

19



Europäisches Patentamt
European Patent Office
Office européen des brevets

11

Publication number

0 141 927
A2

12

EUROPEAN PATENT APPLICATION

21 Application number: 84109539.1

22 Date of filing: 10.08.84

51 Int. Cl.: C 07 D 473/18, C 07 D 473/22,
C 07 D 473/16, C 07 D 473/24,
C 07 D 473/40, C 07 F 9/65
// A61K31/52, C07C31/22,
C07C43/164, C07C69/18,
C07C69/734

30 Priority: 18.08.83 GB 8322199
21.09.83 GB 8325271
30.03.84 GB 8408322

43 Date of publication of application: 22.05.85
Bulletin 85/21

84 Designated Contracting States: BE CH DE FR GB IT LI
NL SE


71 Applicant: BEECHAM GROUP PLC, Beecham House
Great West Road, Brentford Middlesex TW8 9BD (GB)

72 Inventor: Jarvest, Richard Lewis, 29 Beaconsfield Road,
Surliton Surrey (GB)
Inventor: Harnden, Michael Raymond, 47 Guildford
Road, Horsham Sussex (GB)

74 Representative: Russell, Brian John, European Patent
Attorney Beecham Pharmaceuticals Great Burgh Yew
Tree Bottom Road, Epsom Surrey, KT18 5XQ (GB)

54 Antiviral guanine derivatives.

57 A compound of formula (I)



HO-CH₂-CH-CH₂-OH

or a salt or acyl derivative thereof, in which X represents chlorine,
C₁₋₆ alkoxy, phenoxy, phenyl C₁₋₆ alkoxy, NH₂, -OH or -SH, is
useful in treating viral infections.

—REFERENCE BD

ACTORUM AG

EP 0 141 927 A2

This Page Blank (uspto)

19



Europäisches Patentamt
European Patent Office
Office européen des brevets

11

Publication number

0 141 927
A2

12

EUROPEAN PATENT APPLICATION

21 Application number: 84109539.1

22 Date of filing: 10.08.84

51 Int. Cl.: C 07 D 473/18, C 07 D 473/22,
C 07 D 473/16, C 07 D 473/24,
C 07 D 473/40, C 07 F 9/65
// A61K31/52, C07C31/22,
C07C43/164, C07C69/18,
C07C69/734

références, formules, pages à photocopier, etc	No	classement
6. p 0, 18-21, claims	1.	C07D 473/00 B2 -
7. p 0, 54, (ex 35), claims	2.	INF C07D 473/16 -
8. p 25	3.	INF. C07D 473/22 -
9. p 19	4.	INF C07D 473/24 -
10. p 27, 28	5.	INF. C07D 473/40 -
11. p 29	6.	INF. C07D 319/06
12. p. 26	7.	INF C07F 9/65 B53V
	8.	inf 124 B 69/66 C2 B2 c
	9.	inf 124 C07C 31/22
	10.	inf 124 B 69/16
	11.	inf 124 B 69/62 B2
	12.	inf C07C 43/178 C6C

EP 84109539.1.

Berlin

C07C 69/63

C07C 69/16

C07C 69/63



Modèle 350

This Page Blank (uspto)


- 1 -

TITLE AND FIELD
SECTION 1

COMPOUNDS

The present invention relates to compounds having antiviral activity, processes for their preparation and pharmaceutical compositions containing them.

The compound 9-(4-hydroxy-3-hydroxymethylbut-1-yl) guanine of formula (A)




is disclosed in Synthetic Communications, 2(6), 345-351 (1972) but no pharmaceutical activity has been indicated for the compound in this or any other published document. We have repeated the synthesis of the compound as described in the above publication, and have shown that the product is a mixture of the compound of formula (A), its monobenzyl ether and its dibenzyl ether, this mixture having a melting point and uv spectrum in agreement with those reported in the publication for the supposedly 'pure' compound of formula (A). Our analysis of the product produced by the above synthesis showed that it contained 45-50% by weight of the compound of formula (A), 45-50% by weight of the monobenzyl ether and 5% or less by weight of the dibenzyl ether.

- 2 -

By different synthetic routes, we have prepared the compound of formula (A) in a substantially pure form and have found that it has anti-viral activity. This activity is also shown by certain derivatives of the compound of formula (A).

According to the present invention there is provided a compound of formula (I)



or a salt, phosphate ester or acyl derivative thereof, in which X represents chlorine, straight or branched chain C₁-6 alkoxy, preferably methoxy, phenoxy, phenyl C₁-6 alkoxy, -NH₂, -OH or -SH with the proviso that, when X is -OH, the compound of formula (I) is in a purity state of greater than 50% by weight of pure compound.

The term 'acyl derivative' is used herein to include any derivative of the compounds of formula (I) in which one or more acyl groups are present. Such derivatives include biological precursors of the compounds of formula (I) in addition to those derivatives which are per se biologically active.

- 3 -

Examples of acyl derivatives of the compounds of formula (I) are those wherein one or both of the hydrogens in the acyclic OH groups, and/or one of the hydrogen atoms in the -NH_2 group, are replaced by $\text{R}-\text{C}-$ groups, wherein R is hydrogen or an alkyl, aryl,



aralkyl or heterocyclyl group.

Examples of alkyl groups R include straight and branched chain groups containing up to 18 carbon atoms, preferably up to 6 carbon atoms. Particular examples are methyl, ethyl, t-butyl and pentyl.

Examples of aryl groups R include phenyl optionally substituted with up to five preferably up to three groups.


Examples of aralkyl groups R include phenyl- C_{1-6} alkyl groups such as benzyl.

Examples of heterocyclyl groups R include single or fused rings containing one or two hetero-atoms in each ring, selected from oxygen, nitrogen and sulphur.

Examples of phosphate esters of the compounds of formula (I) include those where one or both of the acyclic -OH groups are replaced by $(\text{HO})_2\text{-P-O-}$ groups



or salts thereof, or where the two -OH groups are replaced by a bridging



group


Salts, phosphate esters and acyl derivatives of the compounds of formula (I) are preferably pharmaceutically acceptable, but non-pharmaceutically

- 4 -

acceptable compounds are also within the scope of the present invention, since these are useful as intermediates in the preparation of pharmaceutically acceptable compounds.

The compounds of formula (I) are defined herein as including tautomers of formula (I), wherein the -OH and -SH substituents are replaced by =O and =S substituents respectively.

A particular group of compounds of the invention are those of formula (II)



or pharmaceutically acceptable salts thereof, in which X is as defined in formula (I), and each of R¹, R² and R³ represents hydrogen or an acyl group of formula R⁴-C(=O)-, in which


R⁴ is C₁-18 alkyl or imidazolyl, or R¹ or R² represents a phosphate ester group of formula (HO)₂-P(=O)-, or R¹ and

R² together represent a

bridging group.

- 5 -

Subject to the aforementioned purity proviso in relation to compounds of the invention, a preferred compound of the present invention is the compound of formula (A)



or a salt or acyl derivative thereof.

In a further aspect of the invention there is provided a compound of formula (A) in a purity state of greater than 60% preferably greater than 80% more preferably greater than 90% and particularly preferably more than 95% by weight of pure compound.

In yet a further aspect of the invention, there is provided an isolated, substantially completely pure compound of formula (A), or a pharmaceutically acceptable salt thereof.

The invention also provides a compound of formula (A) in crystalline form having a melting point of 275-277°C.


The compounds of the present invention have antiviral activity, and are potentially useful in the treatment of infections caused by herpes viruses, such as herpes simplex type 1, herpes simplex type 2 and varicella zoster viruses.

- 6 -

Accordingly, the present invention also provides a compound of formula (I) or a pharmaceutically acceptable salt, phosphate ester or acyl derivative thereof, for use as an active therapeutic substance, and in particular for use in the treatment of viral infections. In this aspect of the invention, the compounds of formula (I) are not subject to the aforementioned purity proviso.

Examples of pharmaceutically acceptable salts of the compounds of formula (I) are those formed with organic bases, preferably with amines such as ethanolamines or diamines; and alkali metals, such as sodium and potassium; and acid addition salts formed with a pharmaceutically acceptable acid such as hydrochloric acid, orthophosphoric acid and sulphuric acid.

The compound of formula (A) or a salt thereof may be prepared by converting the group X in a compound of formula (III).



in which X, excluding - OH, is as defined in formula (I); R^a and R^b , which may be the same or different, are each hydrogen or O- protecting groups, preferably acyl groups; and Y is chlorine or -NHR_c, in which R_c is hydrogen or acyl,

- 7 -

to an -OH group by means of hydrolysis, preferably acid hydrolysis, when X is other than NH_2 , or, when X is $-\text{NH}_2$, by means of a deaminase reaction, or when Y is chlorine and X is -OH, converting Y to a $-\text{NH}_2$ group by reaction with ammonia under pressure in accordance with known methods, and subsequently, if desired, converting the compound of formula (A) to a salt thereof by treatment with an acid or base.


Acyl groups R^a , R^b and R^c may be those of formula $\text{R}-\overset{\text{O}}{\underset{\text{O}}{\text{C}}}-$ as hereinbefore defined.

Examples of groups R^a and R^b in formula (III) are acetyl and cyclic acetal such as isopropylidene. R^c is preferably acetyl or hydrogen.

A preferred process for preparing the compound of formula (A) comprises treating a compound of formula (III) in which X is chlorine, Y is $-\text{NH}_2$ and R^a and R^b are each acetyl, with aqueous mineral acid, preferably hydrochloric acid.

Compounds of formulae (III) when each of R^a , R^b and R^c is hydrogen or acyl, are themselves compounds of the invention, having the additional utility as intermediates for the preparation of the compound of formula (A).


In a further aspect of the invention, compounds of formula (I) or acyl derivatives thereof, together with, compounds of formula (III), may be prepared by treating a compound of formula (IV).



(IV)

- 8 -


in which X is as defined in formula (I) and Y is as defined in formula (III), with a compound of formula (V)



in which R^{a} and R^{b} are as defined in formula (III) and Z is a leaving group such as Cl, Br, or I, preferably Br.

Compounds of formula (IV) are either known compounds or can be made from known compounds by known methods.

Compounds of formula (V) in which Z is bromine may be prepared by brominating a compound of formula (VI).




preferably by treatment with carbon tetrabromide and triphenylphosphine in an organic, aprotic solvent, such as dimethylformamide.

Compounds of formula (V) in which Z is Cl or I may be prepared in an analogous manner.

Compounds of formula (VI) in which R^{a} and R^{b} are identical may be prepared according to the following schematic process.

- 9 -



Acyl derivatives of compounds of formula (I) may also be prepared by acylating an optionally protected compound of formula (I) in accordance with conventional acylating processes known in the art, and where necessary, deprotecting the resulting product.

The acylation reaction may be carried out by using an acylating agent containing a $\text{R}-\overset{\text{O}}{\underset{\text{O}}{\text{C}}}-$ group, wherein R is as hereinbefore defined.

In a particular aspect of this process, the acylating agent contains the $\text{R}^4-\overset{\text{O}}{\underset{\text{O}}{\text{C}}}-$ group, in which R^4 is C_1 -18 alkyl, or is N,N'-carbonyldiimidazole.

Examples of acylating agents suitable for the above process are carboxylic acids, acid halides or acid anhydrides, preferably anhydrides or acids.

- 10 -

When the acylating agent is a carboxylic acid, a coupling agent such as dicyclohexylcarbodiimide should be included, but this is not necessary when the acylating agent is an acid anhydride.

The acylation reaction may produce a single acyl derivative of a compound of formula (I), or a mixture of derivatives, depending on a number of factors, such as the relative amounts and chemical natures of reactants, the physical conditions of the reaction, and the solvent system. Any mixture produced in this way may be separated into its pure components using standard chromatographic techniques.

The above described acylation process of the invention can yield mono-, di-, or tri-acylated derivatives of compounds of formula (I) according to the form of protection/deprotection utilised. The following are examples of products obtained by different methods:

- (a) Di-acylated derivatives of the two acyclic-OH groups may be obtained by direct acylation of unprotected compounds of formula (I) or acylation of protected intermediates of compounds of formula (I) in which the -NH_2 group is protected by, for example, a monomethoxytrityl group, and subsequent deprotection by treatment with acid.
- (b) Mono-acylated derivatives of one of the acyclic -OH groups may be obtained by acylation of protected intermediates of compounds of formula (I) in which the -NH_2 group and the other acyclic -OH group are both protected by, for example, monomethoxytrityl groups, and subsequent deprotection by acid treatment.

- 11 -

- (c) Mono-acylated derivatives of the NH_2 group may be obtained by acylation of protected intermediates of compounds of formula (I) in which both acyclic - OH groups are protected by, for example trimethylsilyl groups, and subsequent deprotection.

The various protected intermediates of compounds of formula (I) may be prepared in accordance with standard procedures by, for example, treatment of the compounds with monomethoxytrityl chloride (for processes (a) and (b)) or with chlorotrimethylsilane (for process (c)).


Protected intermediates of compounds of formula (I) may also be used to prepare phosphate esters of the compounds.

Accordingly, in a further process aspect of the invention, there is provided a process for preparing a mono-phosphate ester of a compound of formula (I) which comprises treating a protected intermediate of the compound of formula (I) in which one of the acyclic - OH groups and the -NH_2 group are protected, preferably by monomethoxytrityl groups, with cyano ethyl phosphoric acid and subsequently deprotecting the resultant product by treatment with acid, preferably acetic acid.

If desired, the reaction product after treatment with cyano ethyl phosphoric acid is treated with aqueous ammonia, which yields the ammonium salt of the phosphate ester as the final product.

- 12 -

Compounds of formula (I) or salts thereof may also be prepared by hydrolysing the 1,3-dioxane ring of a compound of formula (VII).



in which X is as defined in formula (I) and R^C is as defined in formula (III), provided that R^C is not acyl when X is other than OH, and subsequently, if desired, converting the compound of formula (I) thus formed to a salt by treatment with an acid or base.

When R^C is an acyl group, a basic N-deprotection step is required to form the compound of formula (A). This can be carried out prior to or after hydrolysis by treatment with, for example, (i) a solution of NaOMe in CH_3OH or (ii) a solution of NH_3 in CH_3OH .

Preferably the hydrolysis of compounds of formula (VII) is carried out in acid medium. The compounds of formula (VII) in which X is alkoxy, phenoxy, phenylalkoxy or -SH are conveniently prepared in situ by reacting the compound of formula (VII) in which X is chlorine with an additional reactant containing an X^1 .

- 13 -

substituent, wherein X^1 is alkoxy, phenoxy, phenylalkoxy or sulphur. These intermediates can then be hydrolysed to compounds of formula (I) without isolating them from the reaction mixture.

The additional reactant containing the X^1 moiety may be a sodium alkoxide, phenoxide or phenylalkoxide, or sodium hydrosulphide (when X^1 is sulphur).

Acid hydrolysis of a compound of formula (VII) in which X is chlorine will yield a compound of formula (I) in which X is chlorine, or a compound of formula (A) depending on acid strength and reaction conditions.


For example, treatment of the compound of formula (VII) in which X is chlorine with dilute HCl (1.0M) at 60°C for 24 hours or with 2 M HCl under reflux for 1.5 hours, will yield the compound of formula (A). Treatment of the same compound of formula (VII) with 2 M HCl in tetrahydrofuran at room temperature will yield the compound of formula (I) in which X is chlorine.

If desired, the compound of formula (VII) in which X is chlorine may be converted to the compound of formula (VII) in which X is amino, prior to acid hydrolysis.


The conversion may be achieved by treatment with sodium azide in dimethylformamide to form an azido intermediate in which X is replaced by an azide moiety, followed by reduction of the intermediate with ammonium formate/palladium-on-charcoal in methanol.

- 14 -

The intermediate compound of formula (VII) in which X is chlorine and R^C is hydrogen may be prepared by treating a compound of formula (VIII).




with a compound of formula (IX)



The reaction may be carried out in an inert organic solvent, preferably dimethylformamide, in the presence of an inorganic base, preferably potassium carbonate.


The compound of formula (VIII) may itself be prepared by brominating a compound of formula (X)



- 15 -

The reaction is preferably carried out by treating the compound of formula (X) with carbon tetrabromide and triphenylphosphine in an organic, aprotic solvent such as dimethylformamide.

The compound of formula (X) may itself be prepared by treating a compound of formula (XI)



with 2,2-dimethoxypropane and p-toluenesulphonic acid in the presence of acetone or tetrahydrofuran.

The compounds of formulae (IX) and (XI) are known compounds or can be prepared from known compounds by known methods.

The compounds of formulae (VII), (VIII) and (X) are novel intermediates and as such form further aspects of the present invention.

A compound of formula (I) or pharmaceutically acceptable salt, phosphate ester or acyl derivative thereof may be formulated for use in a pharmaceutical composition. Accordingly, in a further aspect of the invention, there is provided a pharmaceutical composition which comprises a compound of formula (I) or pharmaceutically acceptable salt, phosphate ester or acyl derivative thereof together with a pharmaceutically acceptable carrier or excipient.

- 16 -

A composition which may be administered by the oral route to humans may be compounded in the form of a syrup, tablet or capsule. When the composition is in the form of a tablet, any pharmaceutical carrier suitable for formulating such solid compositions may be used, for example magnesium stearate, starch, lactose, glucose, rice, flour and chalk. The composition may also be in the form of an ingestible capsule, for example of gelatin, to contain the compound, or in the form of a syrup, a solution or a suspension. Suitable liquid pharmaceutical carriers include ethyl alcohol, glycerine, saline and water to which flavouring or colouring agents may be added to form syrups. The compounds may also be presented with a sterile liquid carrier for injection.

The composition may also be formulated for topical application to the skin or eyes.

For topical application to the skin, the composition may be in the form of a cream, lotion or ointment. These formulations may be conventional formulations well known in the art, for example, as described in standard books of pharmaceuticals and cosmetics, such as Harry's Cosmeticology published by Leonard Hill Books and the British Pharmacopoeia.

The composition for application to the eyes may be a conventional eye-drop composition well known in the art, or an ointment composition.

Preferably, the composition of this invention is in unit dosage form or in some other form that the patient may administer to himself a single dose. A suitable dosage unit might contain from 50 mg to 1 g of active ingredient, for example 100 to 500 mg. Such doses may

- 17 -

be administered 1 to 4 times a day or more usually 2 or 3 times a day. The effective dose of compound will in general be in the range of from 1.0 to 20 mg/kg of body weight per day or more usually 2.0 to 10 mg/kg per day.

In a further aspect of the invention there is provided a method of treating viral infections in a human or non-human animal, which comprises administering to the animal an effective, non-toxic amount of a compound of formula (I) or a pharmaceutically acceptable salt, phosphate ester or acyl derivative thereof.

The preparation of compounds of the invention is illustrated by the following Examples.

Example 15-(2-Hydroxyethyl)-2,2-dimethyl-1,3-dioxan

To a suspension of lithium aluminium hydride (2.87g, 76mmol) in tetrahydrofuran (125ml), a solution of triethyl 1,1,2-ethanetricarboxylate (9.2ml, 9.85g, 40mmol) in tetrahydrofuran (25ml) was added dropwise with stirring over 2 hours. Excess reagent was then quenched with aqueous tetrahydrofuran (1:2). The inorganic salts were filtered off and washed with ethanol (100ml). The filtrate and washings were combined and the solvent was evaporated under reduced pressure to afford a colourless oil (4.85g). To a suspension of this oil in acetone (100ml), 2,2-dimethoxypropane (25ml) and p-toluenesulphonic acid monohydrate (2.3g, 12mmol) were added and the mixture was stirred for 1 hour. The resulting solution was neutralised with Amberlite IR 45 (OH⁻ form, methanol washed), filtered and the solvent evaporated under reduced pressure. The residue was purified by column chromatography on silica gel, eluting with chloroform-methanol mixtures (40:1 and 25:1) to afford 5-(2-hydroxyethyl)-2,2-dimethyl-1,3-dioxan as a colourless liquid (3.01g, 47%); ν_{\max} (film) 3420, 2940, 1375, 1200 and 1080 cm^{-1} ; δ_{H} (CDCl_3) 1.34-1.70 (8H, m, $\text{C}(\text{CH}_3)_2$ and $\text{CH}_2\text{CH}_2\text{OH}$), 1.7-2.1 (1H, m, CH), 2.15 (1H, br, D_2O exchangeable, OH), and 3.5-4.0 (6H, m, 3 x CH_2O); (Found: C, 58.33; H, 10.11%. $\text{C}_8\text{H}_{16}\text{O}_3 \cdot 0.25\text{H}_2\text{O}$ requires C, 58.34; H, 10.10%. $[\text{M}-\text{CH}_3]^+$ found 145.0864; $\text{C}_7\text{H}_{13}\text{O}_3$ requires 145.0865).

Alternative Procedure for Preparation of 5-(2-Hydroxyethyl)-
2,2-dimethyl-1,3-dioxan (Example 1)

A. 1,4-Dihydroxy-2-hydroxymethylbutane

To a refluxing solution of triethyl 1,1,2-ethanetricarboxylate (46ml, 200mmol) and sodium borohydride (20g, 530mmol) in t-butanol (400ml), methanol was added in 3 aliquots over 30 minutes (total 25ml). The solution was refluxed for a further 30 minutes and allowed to cool. Hydrochloric acid (5M) was carefully added to neutralise the solution. The solution was filtered and the inorganic residue was extracted with ethanol (2 x 100ml) and filtered. The organic solutions were combined and the solvent removed. The residue was extracted with ethanol (120ml) and the solution filtered. The solvent was removed to afford 1,4-dihydroxy-2-hydroxymethylbutane (24g, 100%); δ_H (D_2O) 1.53 (2H, q, J 6Hz, 3-H), 1.75 (1H, m, 2-H), 3.57 (4H, d, J 6Hz, 1-H and 1'-H), and 3.64 (2H, t, J 6Hz, 4-H).

B. 5-(2-Hydroxyethyl)-2,2-dimethyl-1,3-dioxan

To a solution of 1,4-dihydroxy-2-hydroxymethylbutane (12g, 100mmol) in tetrahydrofuran (25ml), 2,2-dimethoxypropane (13.5g, 110mmol) and p-toluenesulphonic acid monohydrate (0.57g, 3mmol) were added. The solution was stirred for 0.5 hour at room temperature and was then neutralised by addition of triethylamine. The solvent was removed and the residue purified by column chromatography on silica gel eluting with chloroform-methanol mixtures to afford 5-(2-hydroxyethyl)-2,2-dimethyl-1,3-dioxan as a clear colourless liquid (6.5g, 41%).

Example 25-(2-Bromoethyl)-2,2-dimethyl-1,3-dioxan

To an ice-cooled solution of 5-(2-hydroxyethyl)-2,2-dimethyl-1,3-dioxan (1.92g, 12mmol) and carbon tetrabromide (7.96g, 24mmol) in dimethylformamide (100ml), triphenylphosphine (6.30g, 24mmol) was added and the solution was left at 4°C overnight. To this solution methanol (20ml) was added and the solvent was then evaporated under reduced pressure. The residue was purified by column chromatography on silica gel, eluting with hexane-acetone (12:1) to afford 5-(2-bromoethyl)-2,2-dimethyl-1,3-dioxan as a clear colourless liquid (0.89g, 40%); ν_{max} (film) 2940, 1370, 1270, 1260, 1200, and 1070 cm^{-1} ; δ_{H} (CDCl_3) 1.42 (6H, s, $\text{C}(\text{CH}_3)_2$), 1.94 (3H, m, $\text{CHCH}_2\text{CH}_2\text{Br}$), 3.43 (2H, t, J 7Hz, CH_2Br), and 3.5-4.1 (4H, m, 2 x CH_2O); (Found: C, 42.84; H, 6.93 %. $\text{C}_8\text{H}_{15}\text{BrO}_2$ requires: C, 43.07; H, 6.78 %. $[\text{M}-\text{CH}_3]^+$ found 207.0024; $\text{C}_7\text{H}_{12}\text{BrO}_2$ requires 207.0021).

Alternative Procedure for Preparation of 5-(2-Bromoethyl)-2,2-dimethyl-1,3-dioxan (Example 2)

To an ice-cooled solution of 5-(2-hydroxyethyl)-2,2-dimethyl-1,3-dioxan (6.08g, 38mmol) and carbon tetrabromide (18.90g, 57mmol) in N,N-dimethylformamide (110ml), triphenylphosphine (14.95g, 57mmol) was added. The solution was stirred for 0.5 hour at 0°C. The solution was then diluted with saturated aqueous sodium bicarbonate (55ml) followed by water (55ml), and was extracted with hexane (2 x 150ml). The combined organic layers were dried (magnesium sulphate) and the solvent removed. The residue was placed under high vacuum for 2 hours to remove bromoform. The residue was taken up in a small amount of hexane, filtered and the solvent removed to afford 5-(2-bromoethyl)-2,2-dimethyl-1,3-dioxan (7.40g, 87%) as a colourless oil which crystallised on cooling, m.p. ca. 18°C.

Example 32-Amino-6-chloro-9-[2-(2,2-dimethyl-1,3-dioxan-5-yl)ethyl]-
purine

To a solution of 5-(2-bromoethyl)-2,2-dimethyl-1,3-dioxan (0.75g, 3.7mmol) in dry dimethylformamide (12ml) 2-amino-6-chloropurine (0.68g, 4.0mmol) and then anhydrous potassium carbonate (0.83, 6.0mmol) were added. The solution was stirred at room temperature for 5 hours and left at 4°C overnight. The solution was filtered and the solvent removed. The residue was purified by column chromatography on silica gel, eluting with chloroform-methanol mixtures (80:1 and 60:1) to afford 2-amino-6-chloro-9-[2-(2,2-dimethyl-1,3-dioxan-5-yl)ethyl]purine as a white crystalline solid (0.74g, 64%), m.p. 125-126°C; λ_{max} (H₂O) 223 (ϵ 28,900), 247 (ϵ 5,700), and 310 (ϵ 7,700) nm; ν_{max} (KBr) 3450, 3340, 1635, 1615, 1565, 1470, 1410, and 1375 cm⁻¹; δ_{H} [(CD₃)₂SO] 1.26 (3H, s, CH₃), 1.32 (3H, s, CH₃), 1.45-1.85 (3H, m, CHCH₂CH₂N), 3.51 (2H, dd, J 11Hz and J 7Hz, 2 x H_{ax}), 3.78 (2H, dd, J 11Hz and J 4Hz, 2 x H_{eq}), 4.05 (2H, t, J 7Hz, CH₂N), 6.89 (2H, s, D₂O exchangeable, 2-NH₂), and 8.38 (1H, s, 8-H); (Found: C, 50.37; H, 5.68; N, 22.22 %; M⁺ 311.1136. C₁₃H₁₈ClN₅O₂ requires C, 50.08; H, 5.82; N, 22.46 %; M⁺ 311.1149).

Example 49-(4-Hydroxy-3-hydroxymethylbut-1-yl)guanine

2-Amino-6-chloro-9-[2-(2,2-dimethyl-1,3-dioxan-5-yl)-
ethyl]purine (0.59g, 1.9mmol) in hydrochloric acid (1.0M,
4ml) was stirred at 60°C for 24 hours. The solution was
diluted with water and neutralised with Amberlite IR 45
(OH⁻ form). The mixture was filtered, the resin washed
with water and the solvent evaporated under reduced pressure.
The residue was recrystallised from water to afford 9-(4-
hydroxy-3-hydroxymethylbut-1-yl)guanine (238mg, 49%), m.p.
275-277°C; λ_{max} (H₂O) 253 (ϵ 11,500) and 270 (shoulder,
 ϵ 8,630) nm; ν_{max} (KBr) 3420, 3140, 1690, 1645, and 1605
cm⁻¹; δ_{H} [(CD₃)₂SO] 1.3-1.5 (3H, m, CHCH₂CH₂), 3.42 (4H, d,
J 5Hz, 2 x CH₂O), 3.99 (2H, t, J 7Hz, CH₂N), 4.41 (2H, br,
D₂O exchangeable, 2 x OH), 6.44 (2H, s, D₂O exchangeable,
2-NH₂), 7.71 (1H, s, 8-H), and 10.55 (1H, br, D₂O
exchangeable, 1-H); (M⁺ found 253.1176. C₁₀H₁₅N₅O₃
requires M⁺ 253.1175).

Alternative Procedure for Preparation of 9-(4-Hydroxy-3-hydroxymethylbut-1-yl)guanine (Example 4)

2-Amino-6-chloro-9-[2-(2,2-dimethyl-1,3-dioxan-5-yl)-ethyl]purine (3.74g, 12mmol) in hydrochloric acid (2.0M, 12ml) was heated under reflux for 1.5 hours. The solution was neutralised with aqueous sodium hydroxide (10%) and then allowed to cool. The solution was filtered and the solid washed with water to afford 9-(4-hydroxy-3-hydroxymethylbut-1-yl)guanine as a white crystalline solid (2.18g, 72%), m.p. 275-277°C; (Found: C, 47.31; H, 6.02; N, 27.81 %; $C_{10}H_{15}N_5O_3$ requires C, 47.43; H, 5.97; N, 27.65 %).

Example 5Ethyl 4-benzyloxy-2-ethoxycarbonylbutanoate

69/66 E2B2C

Sodium metal (72g, 3.13mmol) was dissolved in dry ethanol (1.2l) with stirring, then diethyl malonate (477ml, 3.13mmol) and potassium iodide (249.5g, 1.5mmol) were added. An oil containing benzyl-2-bromoethyl ether (278.5g, 1.3mmol) contaminated with ethylene carbonate (198g) was then added slowly to the stirred mixture. On completion of the addition, the reaction mixture was heated under reflux for 18 hours, then poured into ice-water (7.5l) and extracted with ether (3 x 1.6l). The ether fractions were combined, dried (MgSO_4), filtered and evaporated to give an oil (550g). This was vacuum distilled to give ethyl 4-benzyloxy-2-ethoxycarbonylbutanoate (313g, 82%) as a clear oil, b.p. $165-180^\circ/2\text{mm}$. δ_{H} (CDCl_3) 1.22 (6H, t, 2 x CH_3), 2.18 (2H, q, CHCH_2), 3.48 (2H, t, CHCH_2O), 3.55 (1H, t, CH), 4.12 (4H, q, 2 x CO_2CH_2), 4.38 (2H, s, OCH_2Ph), 7.21 (5H, s, Ar).

$$\begin{array}{c} \text{Pr. cat. } \sigma - \text{cat. } \text{CH}_2 \text{ cat. } \text{CH}_2 \text{ CH}_2 \\ | \\ \text{CH}_2 \text{ cat. } \end{array}$$

To a cooled, stirred suspension of lithium aluminium hydride (103g, 2.7mol) in dry ether (2.5l) under nitrogen was added ethyl 4-benzyloxy-2-ethoxycarbonyl-butanoate (362g, 1.23mol) over a period of 3 hours. On completion of the addition, the reaction mixture was allowed to warm to room temperature, and then heated under reflux for 1 hour. It was then re-cooled and the excess lithium aluminium hydride destroyed by dropwise addition of water (100ml), 2M sodium hydroxide (100ml) and water (300ml). The reaction mixture was filtered, and the filter cake washed well with chloroform. The filtrate was dried (MgSO_4), filtered, and evaporated to give 4-benzyloxy-2-hydroxymethylbutan-1-ol as a clear oil (226g, 87%). δ_{H} (CDCl_3) 1.35-2.05 (3H, m, CHCH_2CH_2), 3.30-3.80 (8H, m, 3 x CH_2O , 2 x OH), 4.42 (2H, s, OCH_2Ph), 7.26 (5H, s, Ar).

Example 72-Acetoxymethyl-4-benzyloxybut-1-yl acetate

69/16

To a cooled solution of 4-benzyloxy-2-hydroxymethylbutan-1-ol in dry pyridine (1.1l) was added acetyl chloride (230ml, 3.24mol) over 2 hours, the temperature being maintained below 8°C. On completion of the addition, the reaction mixture was stirred at 5°C for a further 1 hour, then poured into water (4l) and extracted with ethyl acetate (1 x 3l, 1 x 2l). The organic extracts were combined and washed with 2M hydrochloric acid (2 x 1l), water (1l) and brine (1l), dried (MgSO₄), filtered and evaporated to give a pale yellow oil (300g).

Vacuum distillation afforded 2-acetoxymethyl-4-benzyloxybut-1-yl acetate as a colourless oil (220g, 70%) b.p. 160-165°C/0.05mm.

The fraction b.p. 122-160°C/0.05mm (42g) was purified by column chromatography on silica gel, elution with ether-hexane 2:3 affording further 2-acetoxymethyl-4-benzyloxybut-1-yl acetate (27g, 8%). δ_H (CDCl₃) 1.68 (2H, q, CHCH₂CH₂), 2.01 (6H, s, 2 x OCOCH₃), 2.19 (1H, m, CH), 3.50 (2H, t, CH₂OCH₂Ph), 4.03 (4H, d, CH₂OCOCH₃), 4.43 (2H, s, OCH₂Ph), 7.24 (5H, s, Ar).

Example 82-Acetoxymethyl-4-hydroxybut-1-yl acetate

To a solution of 2-acetoxymethyl-4-benzyloxy-but-1-yl acetate (55g, 0.187mol) in ethanol (250ml) was added 10% palladium on carbon (2.5g), and the mixture hydrogenated at atmospheric pressure and room temperature. When the theoretical hydrogen uptake had been achieved (18 hours), the reaction was stopped and filtered through Celite.

Evaporation of the filtrate gave a colourless oil (35g).

This was purified by column chromatography on silica gel, elution with 2% methanol in chloroform affording 2-acetoxymethyl-4-hydroxybut-1-yl acetate as a clear oil (32.9, 86%).

δ_H (CDCl₃) 1.61 (2H, q, CHCH₂CH₂), 2.04 (6H, s, 2 x OCOCH₃), 2.20 (1H, m, CH), 2.61 (1H, br s, D₂O exchangeable, OH), 3.68 (2H, t, CH₂OH), 4.04 (4H, d, 2 x CH₂OCOCH₃).

Example 92-Acetoxymethyl-4-bromobut-1-yl acetate

69/62 B2

A mixture of 2-acetoxymethyl-4-hydroxy-but-1-yl acetate (10g, 49mmol), triphenyl phosphine (19.25g, 73mmol) and carbon tetrabromide (24.4g, 73mmol) was stirred for 18 hours at 4°C in dimethylformamide (150ml). The solvent was then evaporated, and the residue purified by column chromatography on silica gel, eluting with ether-light petroleum 2:3 to afford 2-acetoxymethyl-4-bromobut-1-yl acetate as a pale oil (130g, 99%). δ_H (CDCl₃) 1.73-2.56 (3H, m, CHCH_2CH_2), 2.04 (6H, s, 2 x OCOCH₃), 3.44 (2H, t, CH₂Br), 4.04 (4H, d, 2 x CH₂OCOCH₃).

Examples 11 and 109-(4-Acetoxy-3-acetoxymethylbut-1-yl)-2-amino-6-chloropurine
and 7-(4-Acetoxy-3-acetoxymethylbut-1-yl)-2-amino-6-
chloropurine

A mixture of 2-acetoxymethyl-4-bromobut-1-yl-acetate (13.0g, 48.7mmol), 2-amino-6-chloro-purine (8.25g, 48.7mmol) and anhydrous potassium carbonate (10g, 72.5mmol) was stirred in dry dimethylformamide (100ml) for 18 hours at room temperature. The reaction mixture was then filtered, the filtrate evaporated, and the residue purified by column chromatography on silica gel (500g). Elution with 3% methanol in chloroform afforded 9-(4-acetoxy-3-acetoxymethylbut-1-yl)-2-amino-6-chloropurine as a white solid (13.8g, 80%) mp 135-137°. λ_{\max} (H₂O) 222 (ϵ 28,500), 245 (ϵ 4,800) 307 (ϵ 7,700) nm; ν_{\max} (KBr) 3485, 3310, 3200, 1750, 1730, 1625, 1560, 1525, 1475, 1245 cm⁻¹. δ_{H} [(CD₃)₂SO] (270MHz) 1.85-2.05 (3H, m, CHCH₂CH₂), 2.01 (6H, s, 2 x OCOCH₃), 4.03 (4H, d, 2 x CH₂OCOCH₃), 4.16 (2H, t, CH₂N), 6.88 (2H, br s, D₂O exchangeable, NH₂), 8.17 (1H, s, 8-H). Found C, 47.27; H, 4.94; N, 19.56%. C₁₄H₁₈N₅O₄Cl requires C, 47.26; H, 5.10; N 19.68%.

Subsequent elution with 5% methanol in chloroform afforded 7-(4-acetoxy-3-acetoxymethylbut-1-yl)-2-amino-6-chloropurine as a white solid (2.9g, 17%) mp 174-175° (dec). λ_{\max} (H₂O) 222 (ϵ 25,000), 255 (ϵ 3,900), 318 (ϵ 5,600) nm; ν_{\max} (KBr) 3390, 3310, 3205, 1745, 1735, 1635, 1550, 1505, 1380, 1365, 1310, 1250, 1240 cm⁻¹. δ_{H} [(CD₃)₂SO] (270MHz) 1.86 (2H, q, CHCH₂CH₂), 1.99 (6H, s, 2 x OCOCH₃), 1.95-2.05 (1H, m, CH), 4.03 (4H, d, 2 x CH₂OCOCH₃), 4.38 (2H, t, CH₂N), 6.60 (2H, br s, D₂O exchangeable, NH₂), 8.39 (1H, s, 8-H). Found C, 47.48; H, 5.11; N, 19.52%. C₁₄H₁₈N₅O₄Cl requires C, 47.26; H, 5.10; N, 19.68%.

Alternative procedure for preparation of
9-(4-Hydroxy-3-hydroxymethylbut-1-yl)guanine (Example 4)

A solution of 9-(4-acetoxy-3-acetoxymethyl-but-1-yl)-2-amino-6-chloropurine (15.5g, 43.6mmol) in 2M hydrochloric acid (150ml) was heated under reflux for 2 hours. The solution was then cooled to room temperature and neutralised with 10% sodium hydroxide solution, left to stand at 4°C, and the resulting precipitate filtered off, washed with cold water and recrystallized from water to give 9-(4-hydroxy-3-hydroxymethylbut-1-yl)guanine as a white crystalline solid (9.4g, 85%) mp 275-277°.

Example 129-(4-Hydroxy-3-hydroxymethylbut-1-yl)guanine, sodium salt

To a suspension of 9-(4-hydroxy-3-hydroxymethylbut-1-yl)-guanine (0.30g, 1.2mmol) in water (8ml) was added aqueous sodium hydroxide (1M, 1.2ml). The solvent was removed from the resulting clear solution and trituration with methanol-ethanol afforded 9-(4-hydroxy-3-hydroxymethylbut-1-yl)guanine sodium salt as a white solid (0.32g, 97%); λ_{max} (H_2O , pH7.8) 252 (11,300) nm; ν_{max} (KBr) 3400, 1590, and 1570 cm^{-1} ; δ_{H} [$(\text{CD}_3)_2\text{SO}$] 1.47 (1H, m, 3'-H), 1.70 (2H, q, J 7Hz, 2'-H), 3.3-3.5 (4H, AB part of ABX, 2 x 4'-H), 3.96 (2H, t, J 7Hz, 1'-H), 4.7 (2H, br, D_2O exchangeable, 2 x OH), 5.58 (2H, br.s, D_2O exchangeable, 2-NH₂), and 7.43 (1H, s, 8-H).

Alternative procedure for preparation of
9-(4-Hydroxy-3-hydroxymethylbut-1-yl)guanine (Example 4)

A solution of 9-(4-acetoxy-3-acetoxymethyl-but-1-yl)-2-amino-6-chloropurine (15.5g, 43.6mmol) in 2M hydrochloric acid (150ml) was heated under reflux for 2 hours. The solution was then cooled to room temperature and neutralised with 10% sodium hydroxide solution, left to stand at 4°C, and the resulting precipitate filtered off, washed with cold water and recrystallized from water to give 9-(4-hydroxy-3-hydroxymethylbut-1-yl)guanine as a white crystalline solid (9.4g, 85%) mp 275-277°C.

Example 129-(4-Hydroxy-3-hydroxymethylbut-1-yl)guanine, sodium salt

To a suspension of 9-(4-hydroxy-3-hydroxymethylbut-1-yl)-guanine (0.30g, 1.2mmol) in water (8ml) was added aqueous sodium hydroxide (1M, 1.2ml). The solvent was removed from the resulting clear solution and trituration with methanol-ethanol afforded 9-(4-hydroxy-3-hydroxymethylbut-1-yl)guanine sodium salt as a white solid (0.32g, 97%); λ_{max} (H_2O , pH7.8) 252 (11,300) nm; ν_{max} (KBr) 3400, 1590, and 1570 cm^{-1} ; δ_{H} [$(\text{CD}_3)_2\text{SO}$] 1.47 (1H, m, 3'-H), 1.70 (2H, q, J 7Hz, 2'-H), 3.3-3.5 (4H, AB part of ABX, 2 x 4'-H), 3.96 (2H, t, J 7Hz, 1'-H), 4.7 (2H, br, D_2O exchangeable, 2 x OH), 5.58 (2H, br.s, D_2O exchangeable, 2-NH₂), and 7.43 (1H, s, 8-H).

Example 139-(4-Hydroxy-3-hydroxymethylbut-1-yl)guanine, potassium salt

To a suspension of 9-(4-hydroxy-3-hydroxymethylbut-1-yl)-guanine (0.30g, 1.2mmol) in water (8ml) was added aqueous potassium hydroxide (1M, 1.2ml). The solvent was removed from the resulting clear solution and trituration with methanol-ethanol afforded 9-(4-hydroxy-3-hydroxymethylbut-1-yl)-guanine potassium salt as a white powdery solid (0.34g, 97%); λ_{max} (H_2O , pH 7.4) 252 (12,400); ν_{max} (KBr) 3360, 3180, 1690, 1650, 1585, 1570, and 1480 cm^{-1} ; δ_{H} [$(\text{CD}_3)_2\text{SO}$] 1.47 (1H, m, 3'-H), 1.69 (2H, q, J 7.1Hz, 2'-H), 3.3-3.5 (4H, AB part of ABX, 2 x 4'-H), 3.94 (2H, t, J 7.3Hz, 1'-H), 4.7 (2H, br, D_2O exchangeable, 2 x OH), 5.69 (2H, br.s, D_2O exchangeable, 2-NH₂), and 7.38 (1H, s, 8-H); (Found: C, 41.11; H, 4.91; N, 23.82%; $\text{C}_{10}\text{H}_{14}\text{N}_5\text{O}_3\text{K}$ requires: C, 41.22; H, 4.84; N, 24.04%).

Example 142-Amino-6-chloro-9-(4-hydroxy-3-hydroxymethylbut-1-yl)purine hydrochloride

To a solution of 2-amino-6-chloro-9-[2-(2,2-dimethyl-1,3-dioxan-5-yl)ethyl]purine (0.46g, 1.5mmol) in tetrahydrofuran (4.5ml), hydrochloric acid (2.0M, 0.5ml) was added. A white precipitate formed and after 0.5 hour the solution was diluted with further tetrahydrofuran and was filtered to give 2-amino-6-chloro-9-(4-hydroxy-3-hydroxymethylbut-1-yl)purine hydrochloride (290mg, 63%), decomposed over 165°C; λ_{max} (H₂O, pH5.5) 223 (ϵ 28,400), 245 (ϵ 4,620), and 307 (ϵ 7,620) nm; ν_{max} (KBr) 3370, 3330, 3200, 2500, 1650, 1630, 1595, and 1505 cm⁻¹; δ_{H} [(CD₃)₂SO] 1.53 (1H, m, CHCH₂CH₂), 1.83 (2H, q, J 7Hz, CHCH₂CH₂), 3.35 (4H, d, J 6Hz, 2 x CH₂O), 4.19 (2H, t, J 7Hz, CH₂N), 5.85 (9H, s, D₂O exchangeable, 2 x OH, NH₂, HCl and H₂O), and 8.57 (1H, s, 8-H). (Found: C, 39.04; H, 4.85; N, 22.36 %; C₁₀H₁₄ClN₅O₄·HCl requires C, 38.98; H, 4.91; N, 22.73 %).

Example 152-Amino-9-(4-hydroxy-3-hydroxymethylbut-1-yl)-6-methoxypurine

To a solution of 2-amino-6-chloro-9-[2-(2,2-dimethyl-1,3-dioxan-5-yl)ethyl]purine (0.28g, 0.9mmol) in methanol (2.5ml), methanolic sodium methoxide (1M, 1.0ml) was added and the solution was stirred at 50° for 1.5 hours. The solution was allowed to cool and hydrochloric acid (5M, 0.2ml) and water (0.4ml) were added. After 15 minutes the solution was neutralised with 10% aqueous sodium hydroxide. Silica gel was added and the solvent removed. Column chromatography on silica gel eluting with chloroform-methanol mixtures afforded 2-amino-9-(4-hydroxy-3-hydroxymethylbut-1-yl)-6-methoxypurine (185mg, 77%), m.p. 117-119°C; λ_{max} (H₂O) 213 (ϵ 22,100), 249 (ϵ 6,860), and 280 (ϵ 8,410) nm; ν_{max} (KBr) 3400, 3240, 3210, 1640, 1610, 1590, 1410, and 1395 cm⁻¹; δ_{H} [(CD₃)₂SO] 1.47 (1H, m, CHCH₂CH₂), 1.74 (2H, q, J 7Hz, CHCH₂CH₂), 3.40 (4H, d, J 6Hz, 2 x CH₂O), 3.95 (3H, s, OCH₃), 4.06 (2H, t, J 7Hz, CH₂N), 4.4 (2H, br, D₂O exchangeable, 2 x OH), 6.33 (2H, s, D₂O exchangeable, 2-NH₂), and 7.46 (1H, s, 8-H) (Found: M⁺ 267.1340; C₁₁H₁₇N₅O₃ requires M⁺ 267.1331).

Example 162-Amino-6-ethoxy-9-(4-hydroxy-3-hydroxymethylbut-1-yl)purine

To a suspension of 2-amino-6-chloro-9-[2-(2,2-dimethyl-1,3-dioxan-5-yl)ethyl]purine (0.31g, 1.0mmol) in ethanol (1.5ml) was added sodium ethoxide (1M in ethanol, 1.5ml) and the mixture was stirred at 60° for 1 hour. The resulting solution was allowed to cool, hydrochloric acid (5M, 0.3ml) and water (0.7ml) were added and the solution was stirred for 1 hour at room temperature. The solution was neutralised by addition of aqueous sodium bicarbonate and the solvent was removed. The residue was extracted with chloroform-ethanol (2:1), the solution was filtered and the solvent removed. The residue was purified by column chromatography on silica gel eluting with chloroform-methanol (6:1) to afford 2-amino-6-ethoxy-9-(4-hydroxy-3-hydroxymethylbut-1-yl)purine (0.24g, 85%), m.p. 150-152°C; λ_{max} (H₂O) 213 (24,300), 249 (7,360), and 280 (9,270) nm; ν_{max} (KBr) 3330, 3210, 2900, 1650, 1610, 1580 cm⁻¹; δ_{H} [(CD₃)₂SO] 1.3-1.6 (4H, m, 3'-H and CH₃), 1.73 (2H, q, J 7Hz, 2'-H), 3.2-3.6 (4H, AB part of ABX, 2 x 4'-H), 4.04 (2H, t, J 7Hz, 1'-H), 4.3-4.55 (4H, m, 2H D₂O exchangeable, 2 x OH; D₂O exchange leaves 2H, q, J 7Hz, 6-OCH₂), 6.30 (2H, s, D₂O exchangeable, 2-NH₂), and 7.84 (1H, s, 8-H); (Found: C, 50.91; H, 7.00; N, 24.89%; C₁₂H₁₉N₅O₃ requires C, 51.23; H, 6.81; N, 24.90 %).

Example 172-Amino-6-benzyloxy-9-(4-hydroxy-3-hydroxymethylbut-1-yl)
purine

A suspension of 2-amino-6-chloro-9-[2-(2,2-dimethyl-1,3-dioxan-5-yl)ethyl]purine (0.31g, 1.0mmol) in a solution of sodium benzoate (1M in benzyl alcohol, 2ml) was stirred at 70°C for 1 hour. The resulting solution was allowed to cool, hydrochloric acid (5M, 0.4ml) and water (0.6ml) were added and the solution was stirred for 1 hour at room temperature. The solution was then partitioned between chloroform and water. The aqueous layer was neutralised with aqueous sodium bicarbonate and extracted with chloroform. The combined organic layers were washed with aqueous sodium bicarbonate, dried (magnesium sulphate) and the solvent was removed. The residue was purified by column chromatography on silica gel eluting with chloroform-methanol mixtures (10:1, 5:1) to afford 2-amino-6-benzyloxy-9-(4-hydroxy-3-hydroxymethylbut-1-yl)purine as a white crystalline solid (0.17g, 50%), m.p. 146-147.5°C; λ_{max} (EtOH) 212 (32,300), 250 (8,380), and 283 (10,100) nm; ν_{max} (KBr) 3340, 3220, 1655, 1605, and 1580 cm^{-1} ; δ_{H} [(CD₃)₂SO] 1.3-1.6 (1H, m, 3'-H), 1.72 (2H, q, J 7Hz, 2'-H), 3.38 (4H, AB part of ABX, 2 x 4'-H), 4.03 (2H, t, J 7Hz, 1'-H), 4.36 (2H, t, J 5.5Hz, D₂O exchangeable, 2 x OH), 5.47 (2H, s, PhCH₂), 6.37 (2H, s, D₂O exchangeable, 2-NH₂), 7.3-7.6 (5H, m, C₆H₅), and 7.84 (1H, s, 8-H); (Found: C, 58.89; H, 6.12; N, 19.87%; C₁₇H₂₁N₅O₃ requires C, 59.46; H, 6.16; N, 20.40%).

Example 182-Amino-9-(4-hydroxy-3-hydroxymethylbut-1-yl)-6-thiopurine

A solution of 2-amino-6-chloro-9-[2-(2,2-dimethyl-1,3-dioxan-5-yl)ethyl]purine (0.31g, 1.0mmol) in aqueous sodium hydrosulphide (2M, 3.0ml) and ethanol (1.5ml) was stirred at 70°C for 1 hour. To this solution glacial acetic acid (2.5ml) was added and the mixture was stirred for a further 1 hour at 70°C. The solution was allowed to cool, filtered and the solvent removed. The residue was recrystallised from water to afford 2-amino-9-(4-hydroxy-3-hydroxymethylbut-1-yl)-6-thiopurine (0.13g, 48%), m.p. decomposed at 260°C; δ_H [(CD₃)₂SO] 1.45 (1H, m, CHCH₂CH₂), 1.72 (2H, q, J 7Hz, CHCH₂CH₂), 3.3-3.5 (4H, ABX J_{AB} 10.7Hz, J_{AX} 5.5Hz and J_{BX} 5.8Hz, 2 x CH₂O), 4.02 (2H, t, J 7.4Hz, CH₂N), 4.45 (2H, br, D₂O exchangeable, 2 x OH), 6.77 (2H, s, D₂O exchangeable, 2-NH₂), 7.87 (1H, s, 8-H), and 11.9 (1H, br, D₂O exchangeable, 1-H); λ_{max} (H₂O) 230 (ϵ 16,900), 263 (ϵ 7,210) and 341 (ϵ 25,200) nm; ν_{max} (KBr) 3310, 3130, 1650, 1610, and 1580 cm⁻¹; (Found: C, 44.86; H, 5.60; N, 25.44 %; C₁₀H₁₅N₅O₂S requires C, 44.60; H, 5.61; N, 26.00 %).

Example 172-Amino-6-benzyloxy-9-(4-hydroxy-3-hydroxymethylbut-1-yl)purine

A suspension of 2-amino-6-chloro-9-[2-(2,2-dimethyl-1,3-dioxan-5-yl)ethyl]purine (0.31g, 1.0mmol) in a solution of sodium benzoate (1M in benzyl alcohol, 2ml) was stirred at 70°C for 1 hour. The resulting solution was allowed to cool, hydrochloric acid (5M, 0.4ml) and water (0.6ml) were added and the solution was stirred for 1 hour at room temperature. The solution was then partitioned between chloroform and water. The aqueous layer was neutralised with aqueous sodium bicarbonate and extracted with chloroform. The combined organic layers were washed with aqueous sodium bicarbonate, dried (magnesium sulphate) and the solvent was removed. The residue was purified by column chromatography on silica gel eluting with chloroform-methanol mixtures (10:1, 5:1) to afford 2-amino-6-benzyloxy-9-(4-hydroxy-3-hydroxymethylbut-1-yl)purine as a white crystalline solid (0.17g, 50%), m.p. 146-147.5°C; λ_{max} (EtOH) 212 (32,300), 250 (8,380), and 283 (10,100) nm; ν_{max} (KBr) 3340, 3220, 1655, 1605, and 1580 cm^{-1} ; δ_{H} [(CD₃)₂SO] 1.3-1.6 (1H, m, 3'-H), 1.72 (2H, q, J 7Hz, 2'-H), 3.38 (4H, AB part of ABX, 2 x 4'-H), 4.03 (2H, t, J 7Hz, 1'-H), 4.36 (2H, t, J 5.5Hz, D₂O exchangeable, 2 x OH), 5.47 (2H, s, PhCH₂), 6.37 (2H, s, D₂O exchangeable, 2-NH₂), 7.3-7.6 (5H, m, C₆H₅), and 7.84 (1H, s, 8-H); (Found: C, 58.89; H, 6.12; N, 19.87%; C₁₇H₂₁N₅O₃ requires C, 59.46; H, 6.16; N, 20.40%).

Example 182-Amino-9-(4-hydroxy-3-hydroxymethylbut-1-yl)-6-thiopurine

A solution of 2-amino-6-chloro-9-[2-(2,2-dimethyl-1,3-dioxan-5-yl)ethyl]purine (0.31g, 1.0mmol) in aqueous sodium hydrosulphide (2M, 3.0ml) and ethanol (1.5ml) was stirred at 70°C for 1 hour. To this solution glacial acetic acid (2.5ml) was added and the mixture was stirred for a further 1 hour at 70°C. The solution was allowed to cool, filtered and the solvent removed. The residue was recrystallised from water to afford 2-amino-9-(4-hydroxy-3-hydroxymethylbut-1-yl)-6-thiopurine (0.13g, 48%), m.p. decomposed at 260°C; δ_H [(CD₃)₂SO] 1.45 (1H, m, CHCH₂CH₂), 1.72 (2H, q, J 7Hz, CHCH₂CH₂), 3.3-3.5 (4H, ABX J_{AB} 10.7Hz, J_{AX} 5.5Hz and J_{BX} 5.8Hz, 2 x CH₂O), 4.02 (2H, t, J 7.4Hz, CH₂N), 4.45 (2H, br, D₂O exchangeable, 2 x OH), 6.77 (2H, s, D₂O exchangeable, 2-NH₂), 7.87 (1H, s, 8-H), and 11.9 (1H, br, D₂O exchangeable, 1-H); λ_{max} (H₂O) 230 (ϵ 16,900), 263 (ϵ 7,210) and 341 (ϵ 25,200) nm; ν_{max} (KBr) 3310, 3130, 1650, 1610, and 1580 cm⁻¹; (Found: C, 44.86; H, 5.60; N, 25.44 %; C₁₀H₁₅N₅O₂S requires C, 44.60; H, 5.61; N, 26.00 %).

Example 192-Amino-6-azido-9-[2-(2,2-dimethyl-1,3-dioxan-5-yl)ethyl]-
purine

To a solution of 2-amino-6-chloro-9-[2-(2,2-dimethyl-1,3-dioxan-5-yl)ethyl]purine (0.47g, 1.5mmol) in dry N,N-dimethylformamide (5ml), sodium azide (0.20g, 3.0mmol) was added and the mixture was stirred at 100-110°C for 4 hours. The solvent was removed and the residue washed with water to leave 2-amino-6-azido-9-[2-(2,2-dimethyl-1,3-dioxan-5-yl)ethyl]purine as a crystalline solid (0.36g, 75%), m.p. decomposed at 200°C; λ_{max} (MeOH) 272 (ϵ 8,210) and 301 (ϵ 10,100) nm; ν_{max} (KBr) 1670, 1625, and 1560 cm^{-1} ; δ_{H} (CDCl_3 - CD_3OD) 1.44 (6H, s, $\text{C}(\text{CH}_3)_2$), 1.6-2.2 (3H, m, CHCH_2CH_2), 3.5-3.8 (2H, dd (ABX), J 7Hz and J 11Hz, 2 \times H_{ax}), 3.85-4.15 (2H, dd (ABX), J 4Hz and J 11Hz, 2 \times H_{eq}), 4.29 (2H, t, J 7Hz, CH_2N), and 7.93 (1H, s, 8-H) (Found: C, 48.96; H, 5.66; N, 35.15 %; M^+ 318.1553. $\text{C}_{13}\text{H}_{18}\text{N}_8\text{O}_2$ requires C, 49.05; H, 5.70; N, 35.20 %; M^+ 318.1546).

Example 202,6-Diamino-9-[2-(2,2-dimethyl-1,3-dioxan-5-yl)ethyl]purine

A mixture of 2-amino-6-azido-9-[2-(2,2-dimethyl-1,3-dioxan-5-yl)ethyl]purine (318mg, 1.0mmol), formic acid (0.15ml, 4.0mmol), concentrated ammonia (0.22ml, 4.0mmol), 10% palladium-on-charcoal (30mg) and methanol (10ml) was heated under reflux for 1 hour. The solution was allowed to cool, filtered and the solvent removed. The residue was purified by column chromatography on silica gel eluting with chloroform-methanol mixtures (20:1 and 15:1) to give 2,6-diamino-9-[2-(2,2-dimethyl-1,3-dioxan-5-yl)ethyl]purine (190mg, 65%), m.p. 202-204°C; ν_{\max} (KBr) 1670, 1640, 1595, and 1410 cm^{-1} ; δ_{H} (CDCl_3 - CD_3OD) 1.42 (6H, s, $\text{C}(\text{CH}_3)_2$), 1.6-2.0 (3H, m, CHCH_2CH_2), 3.5-4.2 (6H, m, 2 x CH_2O and CH_2N), and 7.68 (1H, s, 8-H) (Found: C, 52.88; H, 6.78; N, 28.36 %; M^+ 292.1652. $\text{C}_{13}\text{H}_{20}\text{N}_6\text{O}_2$ requires C, 53.41; H, 6.90; N, 28.75 %; M^+ 292.1648).

Example 212,6-Diamino-9-(4-hydroxy-3-hydroxymethylbut-1-yl)purine

A solution of 2,6-diamino-9-[2-(2,2-dimethyl-1,3-dioxan-5-yl)ethyl]purine (180mg, 0.6mmol) in 70% acetic acid (10ml) was stirred for 1 hour at room temperature. The solvent was removed, the residue was suspended in methanol and sodium methoxide was added to neutralise. Column chromatography on silica gel eluting with chloroform-methanol mixtures (6:1, 4:1 and 3:1) gave 2,6-diamino-9-(4-hydroxy-3-hydroxymethylbut-1-yl)purine (95mg, 63%), m.p. 187-190°C; λ_{max} (H_2O , pH6.5) 215 (ϵ 25,500), 255 (ϵ 7,290), and 280 (ϵ 9,170) nm; ν_{max} 3150, 1680, 1650, 1605, 1590, and 1410 cm^{-1} ; δ_{H} [$(\text{CD}_3)_2\text{SO}$] 1.46 (1H, m, CHCH_2CH_2), 1.73 (2H, q, J 7.1Hz, CHCH_2CH_2), 3.3-3.5 (4H, ddd (ABX), J_{AB} 10.6Hz, J_{AX} 5.5Hz and J_{BX} 5.9Hz, 2 x CH_2O), 4.01 (2H, t, J 7.3Hz, CH_2N), 4.41 (2H, br, D_2O exchangeable, 2 x OH), 5.70 (2H, s, D_2O exchangeable, NH_2), 6.56 (2H, s, D_2O exchangeable, NH_2), and 7.69 (1H, s, 8-H) (Found: C, 46.09; H, 6.32; N, 31.69 %; $\text{C}_{10}\text{H}_{16}\text{N}_6\text{O}_2 \cdot 0.1\text{CHCl}_3$ requires C, 45.91; H, 6.14; N, 31.81 %).

Example 229-(4-Acetoxy-3-acetoxymethylbut-1-yl)guanine

A mixture of 9-(4-hydroxy-3-hydroxymethylbut-1-yl)guanine (253mg, 1.0mmol), 4-dimethylaminopyridine (25mg) and acetic anhydride (8.5ml) was stirred for 4 days at room temperature. The acetic anhydride was removed under reduced pressure. The residue was purified by column chromatography on silica gel eluting with chloroform-methanol mixtures (20:1 and 10:1) to afford 9-(4-acetoxy-3-acetoxymethylbut-1-yl)guanine (160mg, 47%) which was recrystallised from methanol, m.p. 202-205°C; ν_{\max} (KBr) 1737, 1690, 1628, 1600, and 1240 cm^{-1} ; δ_{H} [(CD₃)₂SO] 1.79 (2H, q, J 6.7Hz, CHCH₂CH₂), 1.91 (1H, m, CHCH₂CH₂), 2.00 (6H, s, 2 x CH₃), 4.00 (6H, m, 2 x CH₂O and CH₂N), 6.42 (2H, s, D₂O exchangeable, 2-NH₂), 7.71 (1H, s, 8-H), and 10.54 (1H, s, D₂O exchangeable, 1-H); δ_{C} [(CD₃)₂SO] 20.71 (2 x CH₃), 28.29 (C-2'), 34.54 (C-3'), 40.59 (C-1'), 63.58 (2 x C-4'), 116.64 (C-5), 137.58 (C-8), 151.24 (C-4), 153.38 (C-2), 156.87 (C-6), and 170.57 (2 x COO) (Found: C, 49.62; H, 5.70; N, 20.51 %; M^+ 337.1392; C₁₄H₁₉N₅O₅ requires C, 49.85; H, 5.68; N, 20.76 %; M^+ 337.1386).

Examples 23 and 24

9-(4-Propionyloxy-3-propionyloxymethylbut-1-yl)guanine (Example 23) and N²-Propionyl-9-(4-propionyloxy-3-propionyloxymethylbut-1-yl)guanine (Example 24)

A mixture of 9-(4-hydroxy-3-hydroxymethylbut-1-yl)guanine (253mg, 1.0mmol), 4-dimethylaminopyridine (30mg), propionic anhydride (8ml) and N,N-dimethylformamide (15ml) was stirred at room temperature for 66 hours. The solvent was removed and the residue subjected to column chromatography on silica gel eluting with chloroform-methanol mixtures (30:1, 20:1, 10:1). The first compound to elute was N²-propionyl-9-(4-propionyloxy-3-propionyloxymethylbut-1-yl)guanine (200mg, 47%) which was recrystallised from ether-methanol, m.p. 152-154°C; ν_{\max} (KBr) 1740, 1675, 1610, 1560, and 1185 cm^{-1} ; δ_{H} (CDCl_3) 1.14 (6H, t, J 7.5Hz, 2 x $\text{OCOCH}_2\text{CH}_3$), 1.27 (3H, t, J 7.5Hz, $\text{NCOCH}_2\text{CH}_3$), 1.88 (2H, q, J 6.9Hz, CHCH_2CH_2), 2.01 (1H, m, CHCH_2CH_2), 2.35 (2H, q, J 7.5Hz, 2 x $\text{OCOCH}_2\text{CH}_3$), 2.56 (2H, q, J 7.5Hz, $\text{NCOCH}_2\text{CH}_3$), 4.1-4.3 (6H, m, 2 x CH_2O and CH_2N), 7.65 (1H, s, 8-H), 9.14 (1H, s, N-H), and 11.95 (1H, s, N-H) (Found: C, 54.13; H, 6.44; N, 16.19 %; M^+ 421.1958; $\text{C}_{19}\text{H}_{27}\text{N}_5\text{O}_6$ requires C, 54.15; H, 6.46; N, 16.62 %; M^+ 421.1961).

The second compound to elute was 9-(4-propionyloxy-3-propionyloxymethylbut-1-yl)guanine (150mg, 41%) which was recrystallised from methanol, m.p. 204-206°C; ν_{\max} (KBr) 3310, 3150, 1740, 1690, and 1190 cm^{-1} ; δ_{H} [$(\text{CD}_3)_2\text{SO}$] 1.01 (6H, t, J 7.4Hz, 2 x CH_2CH_3), 1.80 (2H, q, J 7.0Hz, CHCH_2CH_2), 1.91 (1H, m, CHCH_2CH_2), 2.29 (4H, q, J 7.5Hz, 2 x CH_2CH_3), 4.01 (6H, m, 2 x CH_2O and CH_2N), 6.37 (2H, s, D_2O exchangeable, 2-NH₂), 7.69 (1H, s, 8-H), and 10.50 (1H, s, D_2O exchangeable, 1-H) (Found: C, 52.28; H, 6.20; N, 18.95 %; $\text{C}_{16}\text{H}_{23}\text{N}_5\text{O}_5$ requires C, 52.59; H, 6.35; N, 19.17 %).

Example 259-(4-Hexanoyloxy-3-hexanoyloxymethylbut-1-yl)guanine

A mixture of 9-(4-hydroxy-3-hydroxymethylbut-1-yl)guanine (253mg, 1.0mmol), dicyclohexylcarbodiimide (0.83mg, 4.0mmol), hexanoic acid (0.38ml, 0.35g, 3.0mmol), 4-dimethylaminopyridine (20mg) and N,N-dimethylformamide (5ml) was stirred for 64 hours at room temperature. The mixture was diluted with water and extracted with chloroform (x 2). The combined organic layers were washed with aqueous sodium bicarbonate, dried (magnesium sulphate) and the solvent removed. The residue was purified by column chromatography eluting with chloroform-methanol mixtures to afford 9-(4-hexanoyloxy-3-hexanoyloxymethylbut-1-yl)guanine (200mg, 45%) which was recrystallised from methanol, m.p. 198.5-201°C; ν_{\max} (KBr) 3340, 3160, 2960, 2930, 1740, 1690, 1650, 1605, and 1170 cm^{-1} ; δ_{H} (CDCl_3) 0.87 (6H, t, J 6.9Hz, 2 x CH_3), 1.28 (8H, m, 2 x $\text{CH}_2\text{CH}_2\text{CH}_3$), 1.60 (4H, quintet, J 7.4Hz, 2 x COCH_2CH_2), 1.90 (2H, q, J 6.9Hz, $\text{CHCH}_2\text{CH}_2\text{N}$), 2.02 (1H, m, $\text{CHCH}_2\text{CH}_2\text{N}$), 2.30 (4H, t, J 7.6Hz, 2 x COCH_2CH_2), 4.13 (6H, m, 2 x CH_2O and CH_2N), 6.42 (2H, s, D_2O exchangeable, 2- NH_2), 7.70 (1H, s, 8-H), and 12.16 (1H, s, D_2O exchangeable, 1-H) (Found: C, 58.97; H, 7.92; N, 15.45 %; $\text{C}_{22}\text{H}_{35}\text{N}_5\text{O}_5$ requires C, 58.78; H, 7.85; N, 15.58 %).

Example 269-(4-Formyloxy-3-formyloxymethylbut-1-yl)guanine

A mixture of 9-(4-hydroxy-3-hydroxymethylbut-1-yl)-guanine (0.23g, 0.9mmol), dicyclohexylcarbodiimide (0.92g, 4.5mmol), formic acid (0.17ml, 4.5mmol), 4-dimethylaminopyridine (20mg) and N,N-dimethylformamide (5ml) was stirred for 40 minutes at room temperature and then quenched by addition of methanol (1ml). The solution was filtered and the solvent removed. The residue was purified by column chromatography on silica gel eluting with chloroform-methanol mixtures (7:1, 4:1) to afford 9-(4-formyloxy-3-formyloxymethylbut-1-yl)guanine which was crystallised from methanol (0.12g, 43%), m.p. 195-198°C; ν_{\max} (KBr) 1720, 1680, 1630, 1600, and 1570 cm^{-1} ; δ_{H} [(CD₃)₂SO] 1.83 (2H, q, J 7.1Hz, 2'-H), 2.01 (1H, m, 3'-H), 4.04 (2H, t, J 7.1Hz, 1'-H), 4.13 (4H, d, J 5.5Hz, 2 x 4'-H), 6.39 (2H, s, D₂O exchangeable, 2-NH₂), 7.70 (1H, s, 8-H), 8.23 (2H, s, 2 x HCOO), and 10.52 (1H, s, D₂O exchangeable, 1-H); (Found: C, 45.40; H, 4.68; N, 21.70%; C₁₂H₁₅N₅O₅ requires: C, 46.60; H, 4.89; N, 22.64%).

Example 279-[4-(N-Imidazolylcarbonyloxy)-3-(N-imidazolylcarbonyloxymethyl)-but-1-yl]guanine

A mixture of 9-(4-hydroxy-3-hydroxymethylbut-1-yl)guanine (253mg, 1.0mmol), N,N'-carbonyldiimidazole (187mg, 1.15mmol), 4-dimethylaminopyridine (20mg) and N,N-dimethylformamide (5ml) was stirred at room temperature. After 2 hours a further quantity of N,N'-carbonyldiimidazole (180mg) was added and stirring was continued for a further 2 hours. The solvent was removed and the residue washed with water, ethyl acetate and hot methanol leaving 9-[4-(N-imidazolylcarbonyloxy)-3-(N-imidazolylcarbonyloxymethyl)but-1-yl]guanine (360mg, 82%), m.p. $>300^{\circ}\text{C}$; ν_{max} (KBr) 3320, 3140, 1760, 1695, 1630, and 1600 cm^{-1} ; δ_{H} [(CD₃)₂SO] 2.0 (2H, q, J 7Hz, CHCH₂CH₂), 2.15-2.40 (1H, m, CHCH₂CH₂), 4.10 (2H, t, J 7Hz, CH₂N), 4.48 (4H, d, J 5Hz, 2 x CH₂O), 6.32 (1H, s, D₂O exchangeable, 2-NH₂), 7.05 (2H, s, imid-H), 7.57 (2H, s, imid-H), 7.75 (1H, s, 8-H), 8.28 (2H, s, imid-H), and 10.53 (1H, s, D₂O exchangeable, 1-H); M/Z 68 (100, imidazole⁺), 44 (70, CO₂⁺), 41 (72, NCHN⁺).

Example 28 & 29

N²-Monomethoxytrityl-9-(4-monomethoxytrityloxy-3-hydroxy-methylbut-1-yl)guanine

and

N²- Monomethoxytrityl -9-(4-hydroxy-3-hydroxymethylbut-1-yl)-guanine

A solution of 9-(4-hydroxy-3-hydroxymethylbut-1-yl)-guanine (4.05g, 16mmol), monomethoxytrityl chloride (10.9g 35mmol), triethylamine (6.7ml) and 4-dimethylaminopyridine (40mg) in N,N-dimethylformamide (50ml) was stirred for 2 hours. The reaction was quenched with methanol and the solvent was removed. The residue was taken up in ethyl acetate and the solution washed with aqueous sodium bicarbonate and water. The solution was dried (magnesium sulphate) and the solvent removed. The residue was purified by column chromatography on silica gel eluting with chloroform-methanol mixtures. The first major product to elute was N²-monomethoxytrityl-9-(4-monomethoxytrityloxy-3-hydroxy-methylbut-1-yl)guanine (4.4g, 34%), m.p. 142-145°C; λ_{\max} (EtOH) 230 (sh. 29,900) and 262 (16,000) nm; ν_{\max} (KBr) 3400, 1680, 1605, 1570, and 1510 cm^{-1} ; δ_{H} [(CD₃)₂SO] 1.24 (2H, m, 2'-H), 1.43 (1H, m, 3'-H), 2.7-2.9 (2H, AB part of ABX, CH₂OC), 3.1-3.4 (2H, AB part of ABX, CH₂OH), 3.42 (2H, t, J 6.7Hz, 1'-H), 3.66 (3H, s, CH₃O), 3.74 (3H, s, CH₃O), 4.35 (1H, t, J 4.8Hz, D₂O exchangeable, OH), 6.7-7.4 (28H, m, Ar-H), 7.44 (1H, s, 8-H), 7.55 (1H, s, D₂O exchangeable, 2-NH), and 10.50 (1H, s, D₂O exchangeable, 1-H); (Found: C, 74.28; H, 5.86; N, 8.64%; C₅₀H₄₇N₅O₅ requires: C, 75.26; H, 5.94; N, 8.78%).

The second major product to elute was N²-monomethoxytrityl-9-(4-hydroxy-3-hydroxymethylbut-1-yl)guanine (1.4g, 17%), m.p. 205-207°C; λ_{\max} (EtOH) 261 (14,500) nm; ν_{\max} (KBr) 3380, 1705, 1680, 1610, 1570, and 1515 cm^{-1} ; δ_{H} [(CD₃)₂SO] 1.25 (3H, m, 2'-H and 3'-H), 3.1-3.3 (4H, m, 2 x 4'-H), 3.52 (2H, t, J 6.6Hz, 1'-H), 3.72 (3H, s, CH₃O), 4.28 (2H, t, J 5.2Hz, D₂O exchangeable, 2 x OH), 6.85-7.35

(14H, m, Ar-H), 7.54 (1H, s, 8-H), 7.56 (1H, s, D₂O exchangeable, 2-NH), and 10.49 (1H, s, D₂O exchangeable, 1-H);
(Found: C, 67.93; H, 6.05; N, 12.90%; C₃₀H₃₁N₅O₄ requires C, 68.55; H, 5.95; N, 13.32%).

Example 309-(4-Pivalyloxy-3-pivalyloxymethylbut-1-yl)guanine

To a solution of N²-monomethoxytrityl-9-(4-hydroxy-3-hydroxymethylbut-1-yl)guanine (0.47g, 0.9mmol) in pyridine (4.5ml) was added pivalyl chloride (0.55ml, 4.5mmol) and the solution was stirred for 45 minutes. The mixture was precipitated in water (45ml) and the resulting precipitate was stirred in 80% acetic acid (10ml) at 80° for 20 minutes. The solvent was removed and the residue was purified by column chromatography on silica gel eluting with chloroform-methanol (10:1) to afford 9-(4-pivalyloxy-3-pivalyloxy-methylbut-1-yl)guanine (0.22g, 58%), m.p. 225-237°C; ν_{max} (KBr) 3430, 2980, 1730, 1690, and 1620 cm^{-1} ; δ_{H} [(CD₃)₂SO] 1.11 (18H, s, 2 x C(CH₃)₃), 1.81 (2H, q, J 6.8Hz, 2'-H), 1.93 (1H, m, 3'-H), 4.0-4.1 (6H, m, 1'-H and 2 x 4'-H), 6.38 (2H, s, 2-NH₂), 7.69 (1H, s, 8-H), and 10.58 (1H, br.s, 1-H); (Found: C, 56.58; H, 7.26; N, 16.14%; C₂₀H₃₁N₅O₅ requires: C, 56.99; H, 7.41; N, 16.62%).

Example 319-(4-Acetoxy-3-hydroxymethylbut-1-yl)guanine

To a solution of N²-monomethoxytrityl-9-(4-monomethoxy-trityloxy-3-hydroxymethylbut-1-yl)guanine (0.72g, 0.9mmol) in pyridine (3ml) was added acetyl chloride (0.21ml, 3.0mmol) and the solution was stirred for 30 minutes. The mixture was precipitated in water (30ml) the resulting precipitate was stirred in 80% acetic acid (10ml) at 80° for 30 minutes. The solvent was removed and the residue purified by column chromatography on silica gel eluting with chloroform-methanol mixtures (7:1, 3:1) to afford 9-(4-acetoxy-3-hydroxymethylbut-1-yl)guanine (0.17g, 64%), m.p. 194-200°C; ν_{max} (KBr) 3330, 3170, 2930, 1730, 1690, 1660, 1610, and 1565 cm^{-1} ; δ_{H} [(CD₃)₂SO] 1.6-1.8 (3H, m, 2'-H and 3'-H), 1.98 (3H, s, CH₃), 3.39 (2H, br, D₂O exchange gives d, J 5Hz, CH₂OH), 3.9-4.1 (4H, m, 1'-H and CH₂OCO), 4.61 (1H, br.t, D₂O exchangeable, OH), 6.44 (2H, s, D₂O exchangeable, 2-NH₂), 7.68 (1H, s, 8-H), and 10.59 (1H, s, D₂O exchangeable, 1-H); (Found: C, 47.91; H, 5.63; N, 21.71%; C₁₂H₁₇N₅O₄ requires: C, 48.81; H, 5.80; N, 23.72%).

Example 329-(4-Benzoyloxy-3-hydroxymethylbut-1-yl)guanine

To a solution of N^2 -monomethoxytrityl-9-(4-monomethoxy-trityloxy-3-hydroxymethylbut-1-yl)guanine (0.72g, 0.9mmol) in pyridine (4ml) was added benzoyl chloride (0.31ml, 2.7mmol) and the solution was stirred for 30 minutes. The mixture was precipitated in water (40ml) and the resulting precipitate was stirred in 80% acetic acid (10ml) at 80° for 45 minutes. The solvent was removed and the residue purified by column chromatography on silica gel eluting with chloroform-methanol mixtures (6:1, 3:1) to afford 9-(4-benzoyloxy-3-hydroxymethylbut-1-yl)guanine (80mg, 25%), m.p. 156-168°C; λ_{max} (MeOH) 231 (15,100) and 254 (13,100) nm; ν_{max} (KBr) 1715, 1690, 1625, and 1600 cm^{-1} ; δ_{H} [(CD_3)₂SO] 1.75-1.90 (3H, m, 2'-H and 3'-H), 3.50 (2H, t, J 5Hz, D_2O exchange leaves d, CH_2OH), 4.07 (2H, t, J 6.9Hz, 1'-H), 4.2-4.35 (2H, AB part of ABX, CH_2OCO), 4.68 (1H, t, J 5.1Hz, D_2O exchangeable, OH), 6.39 (2H, s, D_2O exchangeable, 2-NH₂), 7.5-8.0 (6H, m, C_6H_5 and 8-H), and 10.53 (1H, s, D_2O exchangeable, 1-H); (Found: C, 53.57; H, 5.28; N, 17.95%; $\text{C}_{17}\text{H}_{19}\text{N}_5\text{O}_4 \cdot 0.25 \text{CHCl}_3$ requires: C, 53.51; H, 5.01; N, 18.09%).

Example 339-(4-Hexanoyloxy-3-hydroxymethylbut-1-yl)guanine

To a solution of N²-monomethoxytrityl-9-(4-monomethoxytrityloxy-3-hydroxymethylbut-1-yl)guanine (0.72g, 0.9mmol) in pyridine (4ml) was added hexanoyl chloride (0.38ml, 2.7mmol) and the solution was stirred for 20 minutes. The mixture was precipitated in water (40ml) and the resulting precipitate was stirred in 80% acetic acid (10ml) at 80° for 45 minutes. The solvent was removed and the residue purified by column chromatography on silica gel eluting with chloroform-methanol mixtures (7:1, 5:1) to afford 9-(4-hexanoyloxy-3-hydroxymethylbut-1-yl)guanine (0.12g, 38%), m.p. 179-181°C; ν_{\max} 2960, 2930, 1730, 1690, 1630, and 1600 cm^{-1} ; δ_{H} [(CD₃)₂SO] 0.84 (3H, t, J 6.9Hz, CH₃), 1.24 (4H, m, CH₃(CH₂)₂), 1.50 (2H, quintet, J 7.3Hz, CH₂CH₂CO), 1.6-1.8 (3H, m, 2'-H and 3'-H), 2.26 (2H, t, J 7.3Hz, CH₂CO), 3.40 (2H, t, J 5Hz, D₂O exchange leaves d, CH₂OH), 3.9-4.1 (4H, m, 1'-H and CH₂OCO), 4.60 (1H, t, J 5.1Hz, D₂O exchangeable, 2-NH₂), 7.67 (1H, s, 8-H), and 10.49 (1H, s, D₂O exchangeable, 1-H); (Found: C, 52.63; H, 6.91; N, 18.75%; C₁₆H₂₅N₅O₄ requires: C, 54.69; H, 7.17; N, 19.93%).

Example 349-(4-Hexadecanoyloxy-3-hydroxymethylbut-1-yl)guanine

To a solution of N²-monomethoxytrityl-9-(4-monomethoxy-trityloxy-3-hydroxymethylbut-1-yl)guanine (0.72g, 0.9mmol) in pyridine (4ml) was added hexadecanoyl chloride (0.82ml, 2.7mmol) and the solution was stirred for 30 minutes. The mixture was precipitated in water (40ml) and the resulting precipitate was stirred in 80% acetic acid (8ml) at 80° for 2 hours. The solvent was removed and the residue was purified by column chromatography on silica gel eluting with chloroform-methanol mixtures (10:1, 8:1) to afford 9-(4-hexadecanoyloxy-3-hydroxymethylbut-1-yl)guanine (0.23g, 52%), m.p. 183-191°C; ν_{\max} (KBr) 3340, 3160, 2920, 2850, 1740, 1690, and 1605 cm⁻¹; δ_{H} [(CD₃)₂SO] 0.85 (3H, t, J 6.6Hz, CH₃), 1.23 (24H, m, CH₃(CH₂)₁₂), 1.49 (2H, m, CH₂CH₂CO), 1.6-1.8 (3H, m, 2'-H and 3'-H), 2.26 (2H, t, J 7.3Hz, CH₂CO), 3.39 (2H, t, J 5Hz, D₂O exchange leaves d, CH₂OH), 3.9-4.1 (4H, m, 1'-H and CH₂OCO), 4.60 (1H, t, J 5.2Hz, D₂O exchangeable, OH), 6.38 (2H, s, D₂O exchangeable, 2-NH₂), 7.67 (1H, s, 8-H), and 10.50 (1H, s, D₂O exchangeable, 1-H); (Found: C, 63.83; H, 9.44; N, 14.05%; C₂₆H₄₅N₅O₄ requires: C, 63.51; H, 9.23; N, 14.24%).

Example 359-(4-Hydroxy-3-hydroxymethylbut-1-yl)guanine 4'-phosphate
diammonium salt

To a solution of cyanoethyl phosphoric acid (4.8mmol) in pyridine (6ml) were added N²-monomethoxytrityl-9-(4-monomethoxytrityloxy-3-hydroxymethylbut-1-yl)guanine (1.28g, 1.6mmol) and dicyclohexylcarbodiimide (1.98g, 9.6mmol) and the solution was stirred for 2 hours. Reaction was quenched by addition of water (1ml) and the solvent was removed. To the residue was added concentrated aqueous ammonia and the mixture was stirred at 60° for 3 hours. The solvent was removed and to the residue was added 80% acetic acid (15ml). The mixture was stirred at 80° for 45 minutes and the solvent was removed. The residue was taken up in water (25ml) and extracted with chloroform (4 x 30ml). The aqueous layer was filtered, concentrated and passed down a column of XAD-4 resin, eluting with aqueous methanol mixtures. Fractions containing product were pooled and the solvent removed. The residue was taken up in a small volume of water and the solution was passed through C₁₈-Sep-pak cartridges. Fractions containing product were pooled and the solvent removed to afford 9-(4-hydroxy-3-hydroxymethylbut-1-yl)guanine 4'-phosphate, diammonium salt as a white solid (0.31g, 53%); ν_{\max} (KBr) 3150 (broad), 1690, and 1610 cm^{-1} ; δ_{H} [(CD₃)₂SO/D₂O] 1.57 (1H, m, 3'-H), 1.70 (2H, m, 2'-H), 3.37 (2H, d, J 4.7Hz, CH₂OH), 3.72 (2H, m, CH₂OP), 3.99 (2H, t, J 6.9Hz, 1'-H), and 7.68 (1H, s, 8-H).

Example 36N²-Acetyl-9-(4-hydroxy-3-hydroxymethylbut-1-yl)guanine

To a suspension of 9-(4-hydroxy-3-hydroxymethylbut-1-yl)guanine (0.23g, 0.9mmol) in pyridine (3ml) was added chlorotrimethylsilane (0.25ml, 2.0mmol) and the mixture was stirred for 15 minutes. To this mixture was added acetyl chloride (0.085ml, 1.2mmol) and the mixture was stirred for 30 minutes. Methanol (2ml) was added and the mixture was stirred for a further 30 minutes. The solvent was removed and the residue purified by column chromatography on silica gel eluting with chloroform-methanol mixtures (4:1, 5:2) to give the title compound as its hydrochloride salt. This was dissolved in methanol and stirred with potassium carbonate. The solution was filtered and the solvent removed. The residue was taken up in water and passed through C₁₈-Sep-pak cartridges. Fractions containing product were pooled and the solvent removed to afford N²-acetyl-9-(4-hydroxy-3-hydroxymethylbut-1-yl)guanine (0.11g, 41%), m.p. 143-146°C; λ_{\max} (H₂O) 260 (15,100) nm; ν_{\max} (KBr) 3420, 3200, 2940, 1685, 1615, and 1560 cm⁻¹; δ_{H} [(CD₃)₂SO] 1.46 (1H, m, 3'-H), 1.77 (2H, q, J 7.1Hz, 2'-H), 2.18 (3H, s, CH₃), 3.3-3.5 (4H, m, 2 x 4'-H), 4.13 (2H, t, J 7.4Hz, 1'-H), 4.42 (2H, br.t, J 5Hz, D₂O exchangeable, 2 x OH), 7.98 (1H, s, 8-H), and 11.83 (2H, br, D₂O exchangeable, 1-H and 2-NH).

Example 37N²-Hexanoyl-9-(4-hydroxy-3-hydroxymethylbut-1-yl)guanine

To a suspension of 9-(4-hydroxy-3-hydroxymethylbut-1-yl)-guanine (0.25g, 1.0mmol) in pyridine (5ml) was added chlorotrimethylsilane (0.32ml, 2.5mmol) and the mixture was stirred for 15 minutes. To this mixture was added hexanoyl chloride (0.18ml, 1.3mmol) and the mixture was stirred for 20 minutes. Methanol (2ml) was then added and the mixture was stirred for a further 20 minutes. 1,8-Diazabicyclo[5.4.0]-undec-7-ene (0.57ml, 3.8mmol) and water (0.5ml) were added and the solvent was removed. The residue was purified by column chromatography on silica gel eluting with chloroform-methanol (4:1). Product containing fractions were pooled and the solvent removed. The residue was taken up in water and passed through C₁₈-Sep-pak cartridges to afford N²-hexanoyl-9-(4-hydroxy-3-hydroxymethylbut-1-yl)guanine (50mg, 14%), m.p. 86-88°C; ν_{\max} (KBr) 3400, 2960, 2940, 1675, 1610, and 1560 cm⁻¹; δ_{H} [(CD₃)₂SO] 0.88 (3H, t, J 6.7Hz, CH₃) 1.29 (4H, m, CH₃(CH₂)₂), 1.46 (1H, m, 3'-H), 1.60 (2H, m, CH₂CH₂CO), 1.77 (2H, q, J 7.1Hz, 2'-H), 2.46 (2H, t, J 7.4Hz, CH₂CO) 3.3-3.5 (4H, m, 2 x 4'-H), 4.13 (2H, t, J 7.4Hz, 1'-H), 4.41 (2H, t, J 4.7Hz, D₂O exchangeable, 2 x OH), 7.98 (1H, s, 8-H), 11.65 (1H, br, D₂O exchangeable, NH), and 12.01 (1H, br, D₂O exchangeable, NH); (Found: C, 53.64; H, 7.56; N, 18.95%; C₁₆H₂₅N₅O₄ .0.5H₂O requires: C, 53.32; H, 7.27; N, 19.43%).

a

11.11.13

2-Amino-9-(4-hydroxy-3-hydroxymethylbut-1-yl)-6-isopropoxy-purine

A solution of 2-amino-6-chloro-9-[2-(2,2-dimethyl-1,3-dioxan-5-yl)ethyl]purine (0.25g, 0.8mmol) in isopropanol (2.5ml) containing sodium isopropoxide (0.5M) was stirred at 60° for 25 minutes. After cooling, hydrochloric acid (5M, 0.3ml) and water (0.7ml) were added and the solution was stirred for 15 minutes at room temperature. The solution was neutralised by addition of aqueous sodium bicarbonate and the solvent was removed. The residue was extracted with chloroform-ethanol (2:1) and the solution purified by column chromatography on silica gel eluting with chloroform-methanol (7:1) to afford 2-amino-9-(4-hydroxy-3-hydroxymethylbut-1-yl)-6-isopropoxypurine which was crystallised from chloroform-carbon tetrachloride (0.18g, 76%), m.p. 111.5-113.5°C; δ_H [(CD₃)₂SO] 1.34 (6H, d, J 6.3Hz, C(CH₃)₂), 1.45 (1H, m, 3'-H), 1.74 (2H, q, J 7.2Hz, 2'-H), 3.3-3.5 (4H, AB part of ABX, 2 x 4'-H), 4.06 (2H, t, J 7.3Hz, 1'-H), 4.40 (2H, br, 2 x OH), 5.49 (1H, septet, J 6.3Hz, CH(CH₃)₂), 6.27 (2H, s, 2-NH₂), and 7.83 (1H, s, 8-H).

56b

Example 392-Amino-9-(4-hydroxy-3-hydroxymethylbut-1-yl)-6-phenoxy-purine

To a solution of phenol (113mg, 1.2mmol) in dry dioxan (2.5ml) was added sodium hydride (60% dispersion in oil; 48mg, 1.2mmol). After evolution of hydrogen ceased, 2-amino-6-chloro-9-[2-(2,2-dimethyl-1,3-dioxan-5-yl)ethyl]purine (0.25g, 0.8mmol) was added and the mixture was stirred at 75° for 3.5 hours. After cooling, water (0.8ml) and hydrochloric acid (5M, 0.2ml) were added and the solution was stirred for 30 minutes at room temperature. The solution was neutralised by addition of aqueous sodium bicarbonate and the solvent was removed. The residue was extracted with chloroform-ethanol (2:1) and the solution purified by column chromatography on silica gel eluting with chloroform-methanol (9:1) to afford 2-amino-9-(4-hydroxy-3-hydroxymethylbut-1-yl)-6-phenoxy-purine (145mg, 55%), m.p. 173-175°C; δ_H [(CD₃)₂SO] 1.48 (1H, m, 3'-H), 1.78 (2H, q, J 7.1Hz, 2'-H), 3.3-3.5 (4H, m, 2 x 4'-H), 4.12 (2H, t, J 7.4Hz, 1'-H), 4.42 (2H, t, J 5.1Hz, D₂O exchangeable, 2 x OH), 6.35 (2H, s, D₂O exchangeable, 2-NH₂), 7.2-7.5 (5H, m, C₆H₅), and 7.98 (1H, s, 8-H).

Example of pharmaceutical activityMethod 1

Vero (African Green Monkey Kidney) cells were grown to confluence in 24 well multidishes, each well being 1.6cm in diameter. The cells were infected with Herpes simplex type 1 virus (HFEM strain) and overlaid with 0.5ml of 0.9% agarose (w/v) in maintenance medium. The test compound, prepared in maintenance medium in concentrations ranging from 100 to 0.3 µg/ml in half-log dilution steps, was added in 0.5ml volume. The virus infected cultures were then incubated at 37°C for 4 days before fixing in 4% formaldehyde solution and staining with carbol fuchsin. The dishes were then examined to find what concentration of test compound caused a 50% reduction in the number of virus plaques formed (PDD₅₀ value) and the minimum concentration of test compound which caused cytotoxicity (MTD).

Method 2

MRC-5 cells were infected in suspension with Herpes simplex type 1 virus, strain SC16. The infected cell suspension was dispensed (0.1ml) in 96 well microtitre plates containing the test drugs in maintenance medium in concentrations ranging from 100 to 0.03 µg/ml in half-log dilution steps (0.1ml per well). The plates were then incubated at 37°C for 3 days when the virus cytopathic effect (CPE) in the control wells reached 100%. The plates were fixed in 4% formaldehyde solution and stained with carbol fuchsin. The plates were then examined to find what concentration of test compound reduced the virus CPE by 50% (IC₅₀). Plates using uninfected cells were run in parallel to determine the minimum concentration of test compound which caused cytotoxicity (MTD).

Compounds were also tested against Herpes simplex type 2 virus (MS strain) in Vero cells using Method 1 and in MRC-5 cells using Method 2. In the latter test, the incubation time was reduced to 24 hours.

Results

Example No.	PDD ₅₀ (µg/ml)			
	Herpes simplex type 1 virus		Herpes simplex type 2 virus	
	HFEM strain in Vero cells	SC16 strain in MRC-5 cells	MS strain in Vero cells	MS strain in MRC-5 cells
4	1.3	0.9	2.3	0.6
12	2.2	0.7		
13	1.7	0.7		
15	>100	>100	63	49
22	>100	>100	>100	83
23	>100	25	>100	85
24	>100	>100	>100	65
25	1.9	1.0	1.6	0.9
27	13	1.5	2.8	5.7
31	24	10		
32	95	3		
34	16	2		

None of the compounds was cytotoxic at concentrations up to 100µg/ml in any of the tests.

Method 3

Compounds were administered by oral gavage (0.2mmoles/kg in 0.1ml of 1% carboxymethyl cellulose) to 20g female Balb/C mice which had been starved for 18 hours. Fifteen minutes later, blood was collected from three mice by cardiac puncture using heparinised syringes. Equal aliquots were pooled and an equal volume of 16% trichloroacetic acid added. Following centrifugation (8,500g) to remove precipitated proteins, the resulting mixture was analysed by high performance liquid chromatography using a C₁₈ Nova-Pak cartridge eluted with Buffer A (50mM NaH₂PO₄, pH4.6) and Buffer B (10% Buffer A, 10% water, 80% methanol) in a gradient from 1% to 95% Buffer B. 9-(4-Hydroxy-3-hydroxymethylbut-1-yl) guanine was assayed with a Pye-Unicam PU4021 u.v. detector set at 254nm.


Results

Concentrations of 9-(4-Hydroxy-3-hydroxymethylbut-1-yl)
guanine (Example 4) in the Blood of Mice After Oral
Administration of Derivatives

<u>Administered compound</u>	<u>Concentration of Example 4</u> <u>in Blood (µg/ml)</u>
Example 15	0.2
Example 16	2.3
Example 17	4.8
Example 22	2.0
Example 23	2.5
Example 30	0.3
Example 35	1.1

Claims

1. A compound of formula (I)




or a salt, phosphate ester or acyl derivative thereof, in which X represents chlorine, straight or branched chain C_{1-6} alkoxy, phenoxy, phenyl C_{1-6} alkoxy, $-NH_2$, $-OH$ or $-SH$, with the proviso that, when X is $-OH$, the compound of formula (I) is in a purity state of greater than 50% by weight of pure compound.

2. A compound according to claim 1, in which one or both of the hydrogens in the acyclic $-OH$ groups, and/or one of the hydrogen atoms in the $-NH_2$ group, are replaced by $R-\overset{\overset{O}{\parallel}}{C}-$ groups, wherein R is hydrogen,

alkyl, aryl, aralkyl or heterocyclyl.

3. A compound according to claim 2, in which R is C_{1-6} alkyl, phenyl or benzyl.


4. A compound according to claim 1, of formula (II)



or a pharmaceutically acceptable salt thereof, in which X is as defined in formula (I), and each of R^1 , R^2 and R^3 represents hydrogen or an acyl group of formula

$R^4-\overset{\text{O}}{\underset{\text{O}}{\parallel}}C-$, in which R^4 is C_{1-18} alkyl or imidazolyl, or R^1


or R^2 represents a phosphate ester group of formula $(HO)_2-\overset{\text{O}}{\underset{\text{O}}{\parallel}}P-$, or R^1 and R^2 together represent a



bridging group.

5. A compound according to any one of claims 1 to 4 in which X is -OH, or a tautomer thereof.

6. A compound according to claim 1, of formula (A)



in a purity state of greater than 60% by weight of pure compound, or a pharmaceutically acceptable salt thereof.

7. A compound according to claim 6, of formula (A), in a purity state of greater than 95% by weight of pure compound, or a pharmaceutically acceptable salt thereof.

8. A compound according to claim 6, of formula A, in the form of an isolated, substantially completely pure compound of formula (A), or a pharmaceutically acceptable salt thereof.

9. A compound according to claim 6, of formula (A), in crystalline form having a melting point of 275-277°C.


10. A compound according to any one of claims 1 to 9 without the proviso of claim 1, for use as an active therapeutic substance.

11. A compound according to claim 10 for use in the treatment of viral infections.

12. A pharmaceutical composition comprising a compound according to any one of claims 1 to 9, without the proviso of claim 1, together with a pharmaceutically acceptable carrier or excipient.

- 4 -


13. A process for preparing a compound of formula (A) as defined in claim 6, or a salt thereof, which comprises converting the group X in a compound of formula (III).




in which X, excluding -OH, is as defined in claim 1; R^a and R^b , which may be the same or different, are each hydrogen or O- protecting groups; and Y is chlorine or $-NHR^c$, in which R^c is hydrogen or acyl,

to an -OH group by means of hydrolysis when X is other than NH_2 , or, when X is $-NH_2$ by means of a deaminase reaction or when Y is chlorine, and X is -OH converting Y to a $-NH_2$ group by reaction with ammonia, and subsequently, if desired, converting the compound of formula (A) to a salt thereof by treatment with acid or base.

14. A process for preparing a compound according to claim 1, which comprises treating a compound of formula (IV)



in which X is as defined in formula (I) and Y is as defined in formula (III), with a compound of formula (V)




in which R^a and R^b are as defined in formula (III) and Z is a leaving group.

15. A process for preparing an acyl derivative of a compound of formula (I) as defined in claim 1, which comprises acylating an optionally protected compound of formula (I) and, where necessary, deprotecting the resulting product.


16. A process for preparing a phosphate ester of a compound of formula (I) according to claim 1 which comprises treating a protected intermediate of the compound of formula (I) in which one of the acyclic -OH groups and the -NH₂ group are protected, with cyanoethyl phosphoric acid and subsequently deprotecting the resultant product.

17. A process for preparing a compound of formula (I) as defined in claim 1, or a salt thereof, which comprises hydrolysing the 1,3-dioxane ring of a compound of formula (VII)




in which X is as defined in formula (I) and R^C is as defined in formula (III), provided that R^C is not acyl when X is other than OH, and subsequently, if desired, converting the compound of formula (I) thus formed to a salt by treatment with an acid or base.

18. A compound of formula (VII)




in which X is as defined in claim 1 and R^C is hydrogen or acyl.

19. A compound of formula (VIII)



20. A compound of formula (X)



07D473/22) - (C07D473/24) -
C07D473/40) - (C07D319/06) -
C07F9/65B53Y) - (124B69/66C2B2C) -

07C31/22) - (124B69/16) -
24B69/62B2) - (C07C43/178C6C) -
Office
Office européen des brevets

(11) Publication number:

0 141 927
A3

12-BASIC DOC.-

C07D473/18

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 84109539.1

(22) Date of filing: 10.08.84

(51) Int. Cl.⁴: C 07 D 473/18
C 07 D 473/22, C 07 D 473/16
C 07 D 473/24, C 07 D 473/40
C 07 F 9/65
//A61K31/52, C07C31/22,
C07C43/164, C07C69/18,
C07C69/734

(30) Priority: 18.08.83 GB 8322199
21.09.83 GB 8325271
30.03.84 GB 8408322

(43) Date of publication of application:
22.05.85 Bulletin 85/21

(88) Date of deferred publication of search report: 12.03.86

(84) Designated Contracting States:
BE CH DE FR GB IT LI NL SE

(71) Applicant: BEECHAM GROUP PLC
Beecham House Great West Road
Brentford Middlesex TW8 9BD(GB)


(72) Inventor: Jarvest, Richard Lewis
29 Beaconsfield Road
Surbiton Surrey(GB)

(72) Inventor: Harnden, Michael Raymond
47 Guildford Road
Horsham Sussex(GB)

(74) Representative: Russell, Brian John
European Patent Attorney Beecham Pharmaceuticals
Great Burgh Yew Tree Bottom Road
Epsom Surrey, KT18 5XQ(GB)

(54) Antiviral guanine derivatives.

(57) A compound of formula (I)



or a salt or acyl derivative thereof, in which X represents chlorine, C₁₋₆ alkoxy, phenoxy, phenyl C₁₋₆ alkoxy, NH₂, -OH or -SH, is useful in treating viral infections.

EP 0 141 927 A3

This Page Blank (uspto)

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int Cl 4)
D,A	CHEMICAL ABSTRACTS, vol.78, no.9, March 5, 1973, page 549, ref.no. 58721n; Columbus, Ohio, US U.K. PANDIT et al.: "New class of nucleoside analogs. VIII. Synthesis of N ¹ -pyrimidinyl- and N ⁹ -purinyl- 4'-hydroxy-3-(hydroxymethyl)-buta- nes." & Syn. Commun., 1972, 2(6), 345-51. * Abstract *	1	C 07 D 473/18 C 07 D 473/22 C 07 D 473/16 C 07 D 473/24 C 07 D 473/40 C 07 F 9/65/ A 61 K 31/52 C 07 C 31/22 C 07 C 43/164 C 07 C 69/18 C 07 C 69/734
			TECHNICAL FIELDS SEARCHED (Int Cl 4) C 07 D 473/00 C 07 F 9/00
XXXXXXXXXXXXXXXXXXXXX			
Place of search The Hague		Date of completion of the search 27-08-1985	Examiner LUTEN
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	



CLAIMS INCURRING FEES

The present European patent application comprised at the time of filing more than ten claims.

- ☐ All claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for all claims.
- ☐ Only part of the claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for the first ten claims and for those claims for which claims fees have been paid, namely claims
- ☐ No claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for the first ten claims

X LACK OF UNITY OF INVENTION

The Search Division considers that the present European patent application does not comply with the requirement of unity of invention and relates to several inventions or groups of inventions.

namely: 1) Claims 1-18: Purine compounds, pharmaceutical compositions, preparations
2) Claims 19,20: Compounds VIII and X

- ☐ All further search fees have been paid within the fixed time limit. The present European search report has been drawn up for all claims
- ☐ Only part of the further search fees have been paid within the fixed time limit. The present European search report has been drawn up for those parts of the European patent application which relate to the inventions in respect of which search fees have been paid.

namely claims

- ☒ None of the further search fees has been paid within the fixed time limit. The present European search report has been drawn up for those parts of the European patent application which relate to the invention first mentioned in the claims.

namely claims 1-18